



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

July 28, 2010

MEMORANDUM:

Subject: EPA Reg. No.: 81927-22 Alligare Ecomazapyr 2 SL
DP Barcode: 379826
Case No.: 3078

From: Jolene Trujillo, Biologist *Jolene Trujillo*
Risk Management and Implementation Branch V (7508P)
Pesticide Re-evaluation Division

To: Wilhelmena Livingston, CRM
Risk Management and Implementation Branch V (7508P)
Pesticide Re-evaluation Division

Applicant: Alligare LLC
13 N. 8th Street
Opelika, AL 36801

MJP

FORMULATION FROM EPA Reg. No. 81927-22 LABEL:

	<u>% by wt.</u>
<u>Active Ingredient(s):</u>	
Isopropylamine sale of Imazapyr (2-[4,5-dihydro-4-methyl-4-(1methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid).....	27.8%
<u>Other Ingredient(s):</u>	72.2%
Total	100.0%

BACKGROUND: In the 8 month response to the Imazapyr RED, the registrant has submitted acute toxicity studies to support the reregistration of their product. EPA Reg. No. 81927-22. The MRID's are as follows: [REDACTED]

[REDACTED] The test material used in each of the studies EPA Reg. No. [REDACTED]. All studies were previously reviewed by TRB/RD on [REDACTED] and RMIBV/PRD agrees with this previous review.

RECOMMENDATIONS:

- The acute toxicity studies cited are acceptable to support the reregistration of EPA Reg. No. 81927-22.

The acute toxicity profile for EPA Reg. No. 81927-22 is currently:

Acute Oral	III	Cited
Acute Dermal	III	Cited
Acute Inhalation	IV	Cited
Primary Eye	IV	Cited
Primary Dermal	IV	Cited
Skin Sensitization	non sensitizer	Cited

NOTE: The acute toxicity requirements have been satisfied for the subject product.

LABELING:

ID#: 81927-22 Alligare Ecomazapyr 2 SL

SIGNAL WORD: CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS:

Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wear long sleeved shirt and long pants, socks and shoes and chemical resistant gloves (such as any waterproof material, Category A*).

*If Selection Category A gloves does not provide adequate protection for this product, the registrant should indicate a specific glove category from the EPA chemical resistance chart that will provide adequate protection.

FIRST AID:

IF SWALLOWED: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have product container or label with you when calling a poison control center or doctor or going to treatment. You may also contact 1-800-xxx-xxxx for emergency medical treatment information.

USER SAFETY RECOMMENDATIONS:

User should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

User should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Michael Kellogg
Agent for Alligare, LLC
c/o Pyxis Regulatory Consulting, Inc.
4110 136th St. NW
Gig Harbor, WA 98332

NOV 17 2008

SUBJECT: Application for Pesticide Notification – Clarification of Directions for Use
Alligare Ecomazapyr 2 SL
EPA Reg. No. 81927-22
Application Dated September 23, 2008

Dear Mr. Kellogg:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the label changes requested falls within the scope of PRN 98-10. The label has been date-stamped "Notification" and will be placed in our records.

If you have any questions, please me directly at 703-305-6249 or Terri Stowe of my staff at 703-305-6117.

Sincerely,

A handwritten signature in black ink, appearing to be "L. Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

STG

United States Environmental Protection
Agency Washington, D.C. 20460

OMB Approval 2070-0107
OMB Approval 2070-0057

DATA CALL-IN RESPONSE

INSTRUCTIONS: Please type or print in ink. Please read carefully the attached instructions and supply the information requested on this form.
Use additional sheet(s) if necessary.

1. Company Name and Address ALLIGARE, LLC 4110 136TH ST., NW GIG HARBOR, WA 98332		2. Case # and Name 3021 Arsenal (*) Chemical # and Name 128829 2-(4,5-Dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazole-2-ylidene)-2-propanamine (1:1)		3. Date and Type of DCI and Number 12-Mar-2008 PRODUCT SPECIFIC ID # PDCI-128829-26853	
4. EPA Product Registration	5. I wish to cancel this product registration voluntarily	6. Generic Data		7. Product Specific Data	
		6a. I am claiming a Generic Data Exemption because I obtain the active ingredient from the source EPA registration number listed below.	6b. I agree to satisfy Generic Data requirements as indicated on the attached form entitled "Requirements Status and Registrant's Response."	7a. My product is an MUP and I agree to satisfy the MUP requirements on the attached form entitled "Requirements Status and Registrant's Response."	7b. My product is an EUP and I agree to satisfy the EUP requirements on the attached form entitled "Requirements Status and Registrant's Response."
		N.A.	N.A.		YES
		N.A.	N.A.		YES
		N.A.	N.A.		YES
N.A.	N.A.		YES		
8. Certification I certify that the statements made on this form and all attachments are true, accurate, and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine, imprisonment or both under applicable law. Signature and Title of Company's Authorized Representative <u>DMH Agent</u>		9. Date <u>6/17/08</u>			
10. Name of Company <u>Alligare LLC</u>		11. Phone Number <u>253-553-7369</u>			

PYX REGULATORY CONSULTING INC.

4110 136th St. NW
Gig Harbor, WA 98332

Phone: 253-853-7369
Fax: 253-853-5516
www.PyxisRC.com

November 4, 2008

COURIER DELIVERY

Julia Stokes
Product Reregistration Branch
Special Review and Reregistration Branch (7508P)
Office of Pesticide Programs
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

RE: Arsenal (*) (Case No. 3021; Chemical No. 128829)
Submission of Alligare LLC's 8-Month Response to the Imazapyr Product Specific Data Call-In
Ecomazapyr 2 SL (EPA Reg. No. 81927-22)

Dear Ms. Stokes,


On behalf of Alligare, LLC (Company No. 81927), please find the enclosed 8-month response for the Arsenal (Case No. 3021; Chemical No. 128829) Product Specific Data Call-In for Ecomazapyr 2 SL (EPA Reg. No. 81927-22). In support of this submission, please find the following documents:

1. Application for Reregistration (EPA Form 8570-1)
2. Two copies (2) of the Basic Confidential Statement of Formula dated October 24, 2008
3. Two copies (2) of the Alternate Formulation #1 Confidential Statement of Formula dated October 24, 2008
4. Formulators Exemption (EPA Form 8570-27)
5. Five (5) copies of proposed labeling incorporating changes required by the Imazapyr RED
6. A CD containing an electronic version of the label


Please note that this product is a 100% repack of a currently registered product and the Agency Internal Use Data Matrix, Public file copy of the Data Matrix, and the Certification with Respect to Citation of Data are not required and are not enclosed.

Please feel free to contact me (Janelle@PyxisRC.com; (253) 853-7369) if you have any questions or need any additional information.

Sincerely,


Janelle Kay

Enclosures





United States
Environmental Protection Agency
Washington, DC 20460

☐ Registration
☐ Amendment
☒ Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 81927-22	2. EPA Product Manager J. Stokes	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Alligare LLC/Ecomazapyr 2 SL	PM# SRRD	
5. Name and Address of Applicant (Include ZIP Code) Alligare LLC c/o Pyxis Regulatory Consulting, Inc. 4110 136th St. NW Gig Harbor, WA 98332 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input type="checkbox"/> Notification - Explain below.	<input checked="" type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Arsenal (Case No. 3021; Chemical No. 128829); Imazapyr Product Specific Data Call-In; Submission of 8-Month Response

Section - III

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container

3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container	4. Size(s) Retail Container	5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Janelle Kay	Title Agent	Telephone No. (Include Area Code) (253) 853-7369
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Agent	
4. Typed Name Janelle Kay	5. Date 11/4/08	



United States
Environmental Protection Agency
Washington, DC 20460
Formulator's Exemption Statement
(40 CFR 152.85)

Applicant's Name and Address Alligare, LLC 13 N. 8th Street Opelika, AL 36801	EPA File Symbol/Registration Number 81927-22
	Product Name Alligare Ecomazapyr 2 SL
	Date of Confidential Statement of Formula (EPA Form 8570-4) 10/24/2008

As an authorized representative of the applicant for registration of the product identified above, I certify that:

(1) This product contains the following active ingredient(s):

Imazapyr, isopropylamine salt

(2) Of these, each active ingredient listed in paragraph (4) is present solely as the result of the use of that active ingredient in the manufacturing, formulation or repackaging another product which contains that active ingredient which is registered under FIFRA Section 3, is purchased by us from another person and meets the requirements of 40 CFR section 158.50(e)(2) or (3).

(3) Indicate by checking (A) or (B) below which paragraph applies:

☒ (A) An accurate Confidential Statement of Formula (EPA FORM 8570-4) for the above identified product is attached to this statement. That formula statement indicates, by company name, registration number, and product name, the source of the active ingredient(s) listed in paragraph (1).

OR

☐ (B) The Confidential Statement of Formula (CSF)(EPA Form 8570-4) referenced above and on file with the EPA is complete, current, an accurate and contains the information required on the current CSF.

(4) The following active ingredients in this product qualify for the formulator's exemption.

Product ingredient source information may be entitled to confidential treatment

Source		
Active Ingredient	Product Name	Registration Number
Imazapyr, isopropylamine salt	[REDACTED]	[REDACTED]
Signature 	Name and Title Janelle Kay, Agent	Date 11/4/08



United States
Environmental Protection Agency
Washington, DC 20460

☐ Registration
☐ Amendment
☒ Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 81927-22	2. EPA Product Manager J. Tompkins	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Alligare, LLC / Alligare Ecomazapyr 2 SL	PM# 25	
5. Name and Address of Applicant (Include ZIP Code) Alligare, LLC c/o Pyxis Regulatory Consulting, Inc. 4110 136th St. NW Gig Harbor, WA 98332 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(ii), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

NOTIFICATION
NOV 17 2008

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of a minor label revision per PR Notice 98-10 to clarify the directions for use. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input checked="" type="checkbox"/> Metal	
				<input type="checkbox"/> Plastic	
				<input type="checkbox"/> Glass	
				<input type="checkbox"/> Paper	
				<input type="checkbox"/> Other (Specify) _____	
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per container
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 2.5, 15, 30, 265 gallon		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____			

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			
Name Michael Kellogg	Title Agent	Telephone No. (Include Area Code) (253) 853-7360	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.			6. Date Application Received (Stamped)
2. Signature 	3. Title Agent		
4. Typed Name Michael Kellogg	5. Date 9/23/08		

PYXIS REGULATORY CONSULTING, INC.

4110 136th St. NW
Gig Harbor, WA 98332

Phone: 253-853-7369
Fax: 253-853-5516
www.PyxisRC.com

September 23, 2008

COURIER DELIVERY

Jim Tompkins (PM 25)
Document Processing Desk (**NOTIF**)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 S. Crystal Drive
Arlington, VA 22202

RE: Alligare, LLC – Alligare Ecomazapyr 2 SL (EPA Reg. No. 81927-22)
Notification of Minor Label Revision per PRN 98-10

Dear Mr. Tompkins,


On behalf of Alligare, LLC please find the enclosed notification of a minor label revision for Alligare's Ecomazapyr 2 SL (EPA Reg. No. 81927-22). On page 3 of the enclosed label the word "other" has been removed from the statement "(i.e., DO NOT apply to every other swath)..." so the statement correctly reads, "(i.e., DO NOT apply to every swath)..." The statement is now consistent with the 2nd sentence of the paragraph that reads, "**DO NOT** treat more than one half of the surface area of the water in a single operation and wait at least 10 to 14 days before retreatment."

In support of this notification submission, we submit the following documents:

1. Completed Application for Registration (EPA Form 8570-1)
2. One (1) copy of the Alligare Ecomazapyr 2 SL labeling with changes tracked
3. One (1) copy of the Alligare Ecomazapyr 2 SL labeling with changes incorporated
4. Letter of Authorization

Please feel free to contact me by phone (253) 853-7369 or by email at Mike@PyxisRC.com if you have any questions or need any additional information.


Sincerely,



Michael Kellogg

Enclosures

cc: Jay Golz; Alligare, LLC



Alligare Ecomazapyr 2 SL

Alligare Ecomazapyr 2 SL may only be used for application to control undesirable emergent and floating aquatic vegetation in and around standing and flowing water, including estuarine and marine sites. **Alligare Ecomazapyr 2 SL** may also be used to control undesirable vegetation growing in or around surface water in wetland, riparian and terrestrial habitats where applications may inadvertently contact surface water. **Alligare Ecomazapyr 2 SL** may also be used for cut stump, cut stem and frill and girdle treatments within aquatic sites.

NOTIFICATION

NOV 17 2008

ACTIVE INGREDIENT:

Isopropylamine salt of Imazapyr (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid)*27.8%

OTHER INGREDIENTS72.2%

TOTAL100.0%

*Equivalent to 22.6% 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid or 2 pounds acid equivalent per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCIÓN

PRECAUCION AL USUARIO: Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• DO NOT induce vomiting unless told to so by the poison control center or doctor.• DO NOT give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For medical emergencies involving this product, call 1-800-424-9300.	

EPA Reg. No. 81927-22

Manufactured For:

Alligare, LLC
13 N. 8th Street
Opelika, AL 36801

Net Contents:



PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION! Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing and wash before reuse.

Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistant category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material.
- Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions are given for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

DO NOT apply to water except as specified in this label. Treatment of aquatic weeds may result in oxygen depletion or loss due to decomposition of dead plants. This oxygen loss may cause the suffocation of some aquatic organisms. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

This pesticide is toxic to vascular plants and should be used strictly in accordance with the drift precautions on the label.

PHYSICAL AND CHEMICAL HAZARDS

Spray solutions of **Alligare Ecomazapyr 2 SL** should be mixed, stored and applied only in stainless steel, fiberglass, plastic and plastic-lined steel containers.

DO NOT mix, store or apply **Alligare Ecomazapyr 2 SL** or spray solutions of **Alligare Ecomazapyr 2 SL** in unlined steel (except stainless steel) containers or spray tanks.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Keep containers closed to avoid spills and contamination.

GENERAL INFORMATION

Alligare Ecomazapyr 2 SL is an aqueous solution intended to be mixed with water and surfactant(s) for application to control floating and emergent undesirable vegetation (see **AQUATIC WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL** section and the **ADDITIONAL WEEDS**

CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL section) in or near bodies of water which may be flowing, non-flowing or transient. **Alligare Ecomazapyr 2 SL** may be applied to aquatic sites that include lakes, rivers, streams, ponds, seeps, drainage ditches, canals, reservoirs, swamps, bogs, marshes, estuaries, bays, brackish water, transitional areas between terrestrial and aquatic sites and seasonal wet areas. See **AQUATIC USES, PRECAUTIONS AND RESTRICTIONS** section of this label for precautions, restrictions, and instructions on aquatic uses. **Alligare Ecomazapyr 2 SL** may also be used for cut stump, cut stem and frill and girdle treatments within aquatic sites (see **AERIAL APPLICATIONS AND GROUND APPLICATIONS** sections for additional details).

DO NOT treat more than one half of the surface area of the water in a single operation and wait at least 10 to 14 days between treatments. Begin treatment along the shore and proceed outward in bands to allow aquatic organisms to move into untreated areas.

Read and observe the following directions if aquatic sites are present in terrestrial non-crop areas and are part of the intended treatment area:

Alligare Ecomazapyr 2 SL controls most annual and perennial grasses, broadleaf weeds and many brush and vine species. **Alligare Ecomazapyr 2 SL** will provide some residual control of undesirable species that germinate above the waterline. **Alligare Ecomazapyr 2 SL** controls vegetation by absorption through emergent leaves and stems, from which it is translocated throughout the plant, where it accumulates in rapidly-growing meristematic tissue. Treated plants stop growing soon after treatment. Chlorosis (yellowing of plant tissue) first appears in the newest leaves, and necrosis spreads from this point. In perennials, **Alligare Ecomazapyr 2 SL** is translocated into and kills underground or submerged storage organs to prevent regrowth. Chlorosis and tissue necrosis may not be apparent in some plant species until two or more weeks after application. Complete kill of plants may not occur for several weeks. Applications of **Alligare Ecomazapyr 2 SL** are rainfast within one hour after treatment.

Alligare Ecomazapyr 2 SL will not control plants which are completely submerged or have a majority or their foliage under water.

Alligare Ecomazapyr 2 SL must be applied to the emergent foliage of the target vegetation, since it has little-to-no activity on submerged vegetation. Residual concentrations of **Alligare Ecomazapyr 2 SL** from direct application to water are not expected to be sufficient to provide control of undesirable vegetation. Apply **Alligare Ecomazapyr 2 SL** so as to maximize spray contact onto the target vegetation while minimizing overspray onto adjacent or underlying water. For optimum performance, include a surfactant (see **ADJUVANTS** section for specific recommendations) in the spray solution and apply to actively growing weeds. **Alligare Ecomazapyr 2 SL** may be applied in spot treatments by using low-volume directed application techniques or broadcast by using ground equipment, watercraft or helicopter.

Apply Alligare Ecomazapyr 2 SL with surface or helicopter application equipment in at least 5 gallons of water per acre. When applying by helicopter, follow directions under the **AERIAL APPLICATIONS** section of this label, otherwise refer to section on **GROUND APPLICATIONS** when using surface equipment.

Applications to moving bodies of water should be made while traveling upstream to prevent concentration of this herbicide in water. **DO NOT** apply to bodies of water or portions of bodies of water where emergent and/or floating weeds do not exist.

When target vegetation covers a large percentage of the surface area of impounded water, apply **Alligare Ecomazapyr 2 SL** to alternating strips (i.e., **DO NOT** apply to every swath) to avoid oxygen depletion due to decaying vegetation, since oxygen depletion may result in the suffocation of sensitive aquatic organisms. **DO NOT** treat more than one half of the surface area of the water in a single operation and wait at least 10 to 14 days before retreatment. Begin treatment along the shore and proceed outward in alternating swaths as described above to allow aquatic organisms to move into untreated areas.

Avoid wash-off of sprayed foliage by spray boat or recreational boat backwash for one hour after application.

Apply **Alligare Ecomazapyr 2 SL** at 2 to 6 pints per acre, depending on weed species and weed density. Use higher rates for heavy weed pressure. Refer to the **AQUATIC WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL** section and the **ADDITIONAL WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL** section of this label for specifics. **DO NOT** exceed 6 pints per acre (1.5 lb ai/A) per year.

Alligare Ecomazapyr 2 SL may also be applied as a draw down treatment in areas described above. Apply **Alligare Ecomazapyr 2 SL** to weeds after water has been drained and allow 14 days before reintroduction of water.

PRECAUTIONS FOR AVOIDING INJURY TO NON-TARGET PLANTS

Alligare Ecomazapyr 2 SL can occasionally affect non-target or desirable vegetation by root uptake of the herbicide from treated soil. Injury or loss of non-target plants may result if **Alligare Ecomazapyr 2 SL** is applied onto or near desirable plants, or to areas where their roots extend, or in areas where treated soil may be washed or moved into contact with their root zone.

When making applications along shorelines where desirable plants may be present, exercise caution to avoid spray contact onto their foliage or spray application to the soil in which they are rooted. Shoreline plants with roots that extend into the waters where **Alligare Ecomazapyr 2 SL** has been applied will generally not be adversely affected.

If treated vegetation is to be removed from the application site, **DO NOT** use the vegetative matter as mulch or compost on or around desirable species.

IMPORTANT

DO NOT use on food crops. **DO NOT** apply this product within one-half mile upstream of an active potable water intake in flowing water (i.e., river, stream, etc.) or within one-half mile of an active potable water intake in a standing body of water, such as a lake, pond or reservoir. **DO NOT** apply to water used for irrigation except as described in **APPLICATION TO WATERS USED FOR IRRIGATION** section of this label. Keep away from fertilizers, insecticides, fungicides, and seeds. **DO NOT** drain or flush equipment on or near desirable plants, or onto areas where their roots may extend, or in locations where the chemical or treated soil may be washed or moved into contact with their roots. **DO NOT** use on lawns, walks, driveways, tennis courts or similar areas where roots of desirable vegetation may extend and be exposed to potential injury and/or mortality from root uptake of **Alligare Ecomazapyr 2 SL**. **DO NOT** side trim desirable vegetation with this product unless severe injury or plant death is acceptable. Prevent drift of spray to desirable plants.

Clean application equipment after using this product by thoroughly flushing with water.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Non-crop weed control is not within the scope of the Worker Protection Standard. See the **GENERAL INFORMATION** section of this label for a description of non-crop sites.

DO NOT enter treated areas without protective clothing until sprays have dried.

SPRAY DRIFT MANAGEMENT

Spray Drift: Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determines the potential for spray drift. The applicator and the entity authorizing spraying are responsible for considering all these factors when making decisions.

Spray drift from applying this product may result in damage to sensitive plants adjacent to the treatment area. Only apply this product when the potential for drift to these and other adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, or non-target crops) is minimal. **DO NOT** apply when the following conditions exist that increase the likelihood of spray drift from intended targets: high or gusty winds, high temperatures, low humidity, and temperature inversions.

To minimize spray drift, the applicator should be familiar with and take into account the following drift reduction advisory information. Additional information may be available from state enforcement agencies or the Cooperative Extension on the application of this product.

The best drift management strategy and most effective way to reduce drift potential are to apply large droplets that provide sufficient coverage and control. Applying larger droplets reduces the drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (**see WIND, TEMPERATURE AND HUMIDITY, AND TEMPERATURE INVERSIONS**).

Controlling Droplet Size:

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - **DO NOT** exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of Nozzles - Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is recommended practice. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift. **DO NOT** use nozzles producing a mist droplet spray.

Application Height: Making applications at the lowest possible height (helicopter, ground driven spray bloom) that is safe and practical reduces exposure of droplets to evaporation and wind.

Swath Adjustment: When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the treatment area, the applicator must compensate for this displacement by adjusting the path of the application equipment (e.g. aircraft, ground) upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller droplets, etc.).

Wind: Drift potential is lowest between wind speeds of 3-10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 3 mph due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions: Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud, which can move in unpredictable directions due to the light variable winds common during temperature inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Wind Erosion: Avoid treating powdery dry or light sandy soils when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation.

Aerial Application Methods and Equipment (Helicopters ONLY): Water Volume: Use 2 or more gallons of water per acre. The actual minimum spray volume per acre is determined by the spray equipment used. Use adequate spray volume to produce accurate and uniform distribution of spray particles over the treated area and to avoid spray drift.

Managing Spray Drift from Aerial Applications: Applicators must follow these requirements to avoid off-target drift movement: 1) boom length- the distance of the outermost nozzles on the boom must not exceed $\frac{1}{4}$ the length of the rotor, 2) nozzle orientation- nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees, and 3) application height- without compromising helicopter safety, applications should be made at a height of 10 feet or less above the crop canopy or tallest plants. Applicators must follow the most restrictive use cautions to avoid drift hazards, including those found in this labeling as well as applicable state and local regulations and ordinances.

Ground Application (Broadcast): Water Volume: Use 5 or more gallons of water per acre. The actual minimum spray volume per acre is determined by the spray equipment used. Use adequate spray volume to provide accurate and uniform distribution of spray particles over the treated area and to avoid spray drift.

AQUATIC SPECIES CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL

Alligare Ecomazapyr 2 SL will control the following target species when used as specified in the table below. Rate instructions are expressed as product volume for broadcast applications and as a % solution for directed applications including spot treatments. For % solution applications, **DO NOT** apply more than the equivalent of 3 quarts of Alligare Ecomazapyr 2 SL per acre.

COMMON NAME	SCIENTIFIC NAME	INSTRUCTIONS
Floating Species		
*Duckweed	<i>Lemna minor</i>	2-3 pints/acre (1% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
*Duckweed, Giant	<i>Spirodela polyrriza</i>	2-3 pints/acre (1% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
*Frogbit	<i>Limnobium spongia</i>	1-2 pints/acre (0.5% solution) in 100 GPA water.

COMMON NAME	SCIENTIFIC NAME	INSTRUCTIONS
*Spatterdock	<i>Nuphar luteum</i>	Completely cover all actively-growing emergent foliage. Apply a tank-mix of 2-4 pints/acre Alligare Ecomazapyr 2 SL plus 4-6 pints/acre glyphosate (0.5% Alligare Ecomazapyr 2 SL plus 1.5% glyphosate) in 100 GPA water for best control. Completely cover all actively-growing emergent foliage.
*Water Hyacinth	<i>Eichhornia crassipes</i>	1-2 pints/acre (0.5% solution) in 100 GPA water to actively growing foliage.
*Water Lettuce	<i>Pistia stratiotes</i>	1-2 pints/acre (0.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.

Emerged Species

*Alligatorweed	<i>Alternanthera philoxeroides</i>	1-4 pints/acre (0.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage. DO NOT apply as a tank-mix with glyphosate, since doing so may reduce control.
*Arrowhead, Duck-potato	<i>Sagittaria</i> spp.	1-2 pints/acre (0.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
*Bacopa, lemon	<i>Bacopa</i> spp.	1-2 pints/acre (0.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
*Parrot feather	<i>Myriophyllum aquaticum</i>	Apply 2-4 pints in 100 GPA water to actively growing emergent foliage. Foliage must be above water for sufficient uptake.
*Pennywort	<i>Hydrocotyle</i> spp.	1-2 pints/acre (0.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
*Pickerelweed	<i>Pontederia cordata</i>	2-3 pints/acre (1% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
*Taro, wild; Dasheen; Elephant's Ear; Coco Yam	<i>Colocasia esculentum</i>	4-6 pints/acre (1.5% solution) in 100 GPA with a high quality 'sticker' adjuvant. Completely cover all actively-growing emergent foliage.
*Water lily	<i>Nymphaea odorata</i>	2-3 pints/acre (1% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
*Water Primrose	<i>Ludwigia uruguayensis</i>	4-6 pints/acre (1.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage. DO NOT apply as a tank-mix with glyphosate, since doing so may reduce control.

Terrestrial/Marginal

*Soda Apple, aquatic; Nightshade	<i>Solanum tampicense</i>	2 pints/acre applied to foliage.
*Bamboo, Japanese	<i>Phyllostachys</i> spp.	3-4 pints/acre applied to actively-growing foliage before plants set seed heads. The greater the amount of foliage present at treatment, the greater the exposure to the herbicide for plant uptake, which will result in greater root kill.
Brazilian Pepper;	<i>Schinus terebinthifolius</i>	2-4 pints/acre applied to foliage.

COMMON NAME	SCIENTIFIC NAME	INSTRUCTIONS
Christmasberry Cattail	<i>Typha</i> spp.	2-4 pints (1% solution) applied to actively-growing, green foliage after full leaf elongation. Lower rates will control cattail in the north; higher rates are needed in the south.
Chinese Tallow Tree Cogon grass	<i>Sapium sebiferum</i> <i>Imperata cylindrica</i>	16-24 oz applied to foliage. Burn foliage, till area, then treat in fall at 2 quarts/acre Alligare Ecomazapyr 2 SL plus MSO applied to new growth.
Cordgrass, prairie *Cutgrass *Elephant Grass; Napier Grass *Flowering rush Giant Reed, Wild Cane	<i>Spartina</i> spp. <i>Zizaniopsis miliacea</i> <i>Pennisetum purpureum</i> <i>Butumu typla</i> <i>Arundo donax</i>	4-6 pints applied to actively growing foliage. 4-6 pints applied to actively growing foliage. 3 pints/acre applied to actively growing foliage.
*Golden Bamboo	<i>Phyllostachys aurea</i>	2-3 pints applied to actively growing foliage. 4-6 pints/acre applied in spring to actively growing foliage.
Junglerice Knapweeds	<i>Echinolchloa colonum</i> <i>Centaurea species</i>	3-4 pints/acre applied to actively-growing foliage before plants set seed heads. The greater the amount of foliage present at treatment, the greater the exposure to the herbicide for plant uptake, which will result in greater root kill. 3-4 pints applied to actively growing foliage.
Knotweed, Japanese (see Fallopia japonica) Melaleuca; Paperbark Tree	<i>Polygonum cuspidatum</i> <i>Melaleuca quinquenervia</i>	Russian Knapweed- Apply 2-3 pints plus 1 quart/acre MSO in the fall after senescence begins. 3-4 pints/acre applied postemergence to actively growing foliage. For established stands, apply 6 pints/acre Alligare Ecomazapyr 2 SL plus 6 pints/acre glyphosate plus adjuvant. For best results, use 4 quarts/A methylated seed oil for adjuvant. For ground foliar application, uniformly apply to completely cover all actively-growing foliage. For aerial foliar application, apply in at least two passes in a cross pattern at 10 GPA. For spot treatment, apply a 25% solution of Alligare Ecomazapyr 2 SL plus 25% solution of glyphosate plus 1.25% MSO in water as a frill or stump treatment.
*Nutgrass; Kili'p'opu	<i>Cyperus rotundus</i>	2 pints Alligare Ecomazapyr 2 SL plus 1 quart/acre MSO applied early postemergence.
*Nutsedge	<i>Cyperus</i> spp.	2-3 pints postemergence to foliage or pre-emergence incorporated. Non-incorporated preemergence applications will not provide control.
Phragmites; Common Reed	<i>Phragmites australis</i>	4-6 pints/acre applied to actively-growing, green foliage after leaf elongation. Completely cover all actively-growing foliage. If stand has a substantial amount of old stem tissue, mow or burn, allow to regrow to approximately 5' tall before treatment. Lower rates will control phragmites in the north; higher rates are needed in the south.
*Poison Hemlock	<i>Conium maculatum</i>	2 pints Alligare Ecomazapyr 2 SL plus 1 quart/acre MSO applied preemergence or early

COMMON NAME	SCIENTIFIC NAME	INSTRUCTIONS
Purple Loosestrife	<i>Lythrum salicaria</i>	postemergence up to rosette stage before flowering.
Reed canarygrass	<i>Phalaris arundinacea</i>	1 pint/acre applied to actively growing foliage.
Rose, swamp	<i>Rosa palustris</i>	3-4 pints/acre applied to actively growing foliage.
Russian-Olive	<i>Elaeagnus angustifolia</i>	2-3 pints/ acre applied to actively growing foliage.
Saltcedar, Tamarisk	<i>Tamarix species</i>	2-4 pints/acre or 1% solution applied to foliage. Aerially apply 2 quarts Alligare Ecomazapyr 2 SL plus 0.25% v/v NIS to actively growing foliage during flowering. For spot spraying, use a 1% solution of Alligare Ecomazapyr 2 SL plus 0.25% v/v NIS and spray to wet foliage. After application wait at least two years before disturbing treated saltcedar. Earlier disturbance can reduce overall control.
Smartweed	<i>Polygonum spp.</i>	2 pints/acre applied early postemergence.
Sumac	<i>Rhus spp.</i>	2-3 pints/acre applied to foliage.
Swamp Morning Glory;	<i>Ipomoea aquatica</i>	1-2 pints/acre Alligare Ecomazapyr 2 SL plus
Water Spinach;		1 quart/acre MSO applied at early
Kangkong		postemergence.
Torpedo grass	<i>Panicum repens</i>	4 pints/acre (1-1.5% solution). Completely cover all actively growing foliage.
*White Top; Hoary	<i>Cardaria draba</i>	1-2 pints/acre applied in spring to foliage during flowering.
Cress		2-3 pints/acre Alligare Ecomazapyr 2 SL .
Willow	<i>Salix spp.</i>	Completely cover all actively growing foliage.

* Not approved for use in California

ADDITIONAL WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL

When used as recommended on terrestrial sites, **Alligare Ecomazapyr 2 SL** will provide preemergence or postemergence control with residual control of the following vegetation species. Annual weeds may be controlled by preemergence or postemergence applications of **Alligare Ecomazapyr 2 SL**. For established biennials and perennial vegetation control, postemergence treatments of **Alligare Ecomazapyr 2 SL** are recommended.

The tables below show broadcast rates and indicate relative weed sensitivity. It is important to consider relative weed sensitivity when preparing low volume spray solutions (see **LOW VOLUME** section of **GROUND APPLICATIONS** section), since low volume treatments apply less **Alligare Ecomazapyr 2 SL** per acre than is shown for the broadcast treatments.

The relative sensitivity of the species listed below can also be used to determine the relative risk of causing non-target plant injury if any of those species are considered to be desirable within the treatment area.

Resistant Biotypes: Some weeds listed below may have naturally-occurring biotypes (plants within a given species that have a slightly different but distinct genetic makeup from other plants of that species) that are not effectively controlled by this and/or other herbicides (Oust®) with the ALS/AHAS enzyme-inhibiting mode of action. If naturally-occurring ALS/AHAS-resistant biotypes are present in area, **Alligare Ecomazapyr 2 SL** should be tank-mixed or applied sequentially with a registered herbicide that depends on a different mode of action to ensure control.

GRASSES

COMMON NAME	SPECIES	GROWTH HABIT ²
Apply 2-3 pints per acre¹		
Annual bluegrass	<i>Poa annua</i>	A
Broadleaf signalgrass	<i>Brachiaria platyphylla</i>	A
Canada bluegrass	<i>Poa compressa</i>	P
Downy brome	<i>Bromus tectorum</i>	A
Fescue	<i>Festuca</i> spp.	A/P
Foxtail	<i>Setaria</i> spp.	A
Italian ryegrass	<i>Lolium multiflorum</i>	A
Johnsongrass	<i>Sorghum halepense</i>	P
Kentucky bluegrass	<i>Poa pratensis</i>	P
Lovegrass	<i>Eragrostis</i> spp.	A/P
*Napier grass	<i>Pennisetum purpureum</i>	P
Orchardgrass	<i>Dactylis glomerata</i>	P
Paragrass	<i>Brachiaria mutica</i>	P
Quackgrass	<i>Agropyron repens</i>	P
Sandbur	<i>Cenchrus</i> spp.	A
Sand dropseed	<i>Sporobolus cryptandrus</i>	P
Smooth brome	<i>Bromus inermis</i>	P
Vaseygrass	<i>Paspalum urvillei</i>	P
Wild Oats	<i>Avena fatua</i>	A
Witchgrass	<i>Panicum capillare</i>	A
Apply 3-4 pints per acre¹		
Barnyardgrass	<i>Echinochloa crus-gali</i>	A
Beardgrass	<i>Andropogon</i> spp.	P
Bluegrass, Annual	<i>Poa annua</i>	A
*Bulrush	<i>Scirpus validus</i>	P
Cheat	<i>Bromus secalinus</i>	A
Crabgrass	<i>Digitaria</i> spp.	A
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	A
Fall Panicum	<i>Panicum dichotomiflorum</i>	A
Goosegrass	<i>Eleusine indica</i>	A
Itchgrass	<i>Rottboellia exaltata</i>	A
Lovegrass	<i>Eragrostis</i> spp.	A
*Maidencane	<i>Panicum hemitomon</i>	A
Panicum, Browntop	<i>Panicum fasciculatum</i>	A
Panicum, Texas	<i>Panicum texanum</i>	A
Prairie threeawn	<i>Aristida oligantha</i>	P
Sandbur, Field	<i>Cenchrus incertus</i>	A
Signalgrass	<i>Brachiaria platyphylla</i>	A
Wild barley	<i>Hordeum</i> spp.	A
Wooly Cupgrass	<i>Eriochloa villosa</i>	A
Apply 4-6 pints per acre¹		
Bahiagrass	<i>Paspalum notatum</i>	P
Bermudagrass ³	<i>Cynodon dactylon</i>	P
Big bluestem	<i>Andropogon gerardii</i>	P
Dallisgrass	<i>Paspalum dilatatum</i>	P
Feathertop	<i>Pennisetum villosum</i>	P
Guineagrass	<i>Panicum maximum</i>	P
Saltgrass ³	<i>Distichlis stricta</i>	P
Sand dropseed	<i>Sporobolus cryptandrus</i>	P

COMMON NAME	SPECIES	GROWTH HABIT ²
Sprangletop	<i>Leptochloa</i> spp.	A
Timothy	<i>Phleum pretense</i>	P
Wirestem muhly	<i>Muhlenbergia frondosa</i>	P

BROADLEAF WEEDS

Apply 2-3 pints per acre¹

Burdock	<i>Arctium</i> spp.	B
Carpetweed	<i>Mollugo verticillata</i>	A
Carolina geranium	<i>Geranium carolinianum</i>	A
Clover	<i>Trifolium</i> spp.	A/P
Common chickweed	<i>Stellaria media</i>	A
Common ragweed	<i>Ambrosia artemisiifolia</i>	A
Dandelion	<i>Taraxacum officinale</i>	P
Dog fennel	<i>Eupatorium capillifolium</i>	A
Filaree	<i>Erodium</i> spp.	A
Fleabane	<i>Erigeron</i> spp.	A
Hoary vervain	<i>Verbena stricta</i>	P
Indian mustard	<i>Brassica juncea</i>	A
Kochia	<i>Kochia scoparia</i>	A
Lambsquarters	<i>Chenopodium album</i>	A
*Lespedeza	<i>Lespedeza</i> spp.	P
Miners lettuce	<i>Montia perfoliata</i>	A
Mullein	<i>Verbascum</i> spp.	B
Nettleleaf goosefoot	<i>Chenopodium murale</i>	A
Oxeye daisy	<i>Chrysanthemum leucanthemum</i>	P
Pepperweed	<i>Lepidium</i> spp.	A
Pigweed	<i>Amaranthus</i> spp.	A
Puncturevine	<i>Tribulus terrestris</i>	A
Russian thistle	<i>Salsola kali</i>	A
Smartweed	<i>Polygonum</i> spp.	A/P
Sorrell	<i>Rumex</i> spp.	P
Sunflower	<i>Helianthus</i> spp.	A
Sweet clover	<i>Melilotus</i> spp.	A/B
Tansymustard	<i>Descurainia pinnata</i>	A
Western ragweed	<i>Ambrosia psilostachya</i>	P
Wild carrot	<i>Daucus carota</i>	B
Wild lettuce	<i>Lactuca</i> spp.	A/B
Wild parsnip	<i>Pastinaca sativa</i>	B
Wild turnip	<i>Brassica campestris</i>	B
Woollyleaf bursage	<i>Franseria tomentosa</i>	P
Yellow woodsorrel	<i>Oxalis stricta</i>	P

Apply 3-4 pints per acre¹

Broom snakeweed ⁴	<i>Gutierrezia sarothrae</i>	P
Bull thistle	<i>Cirsium vulgare</i>	B
Burclover	<i>Medicago</i> spp.	A
Chickweed, Mouseear	<i>Cerastium vulgatum</i>	A
Clover, Hop	<i>Trifolium procumbens</i>	A
Cocklebur	<i>Xanthium strumarium</i>	A
Cudweed	<i>Gnaphalium</i> spp.	A

COMMON NAME	SPECIES	GROWTH HABIT ²
Desert Camelthorn	<i>Alhagi pseudalhagi</i>	P
Dock	<i>Rumex</i> spp.	P
Fiddleneck	<i>Amsinckia intermedia</i>	A
Goldenrod	<i>Solidago</i> spp.	P
Henbit	<i>Lamium aplexicaule</i>	A
Knotweed, prostrate	<i>Polygonum aviculare</i>	A/P
Pokeweed	<i>Phytolacca americana</i>	P
Purslane	<i>Portulaca</i> spp.	A
Pusley, Florida	<i>Richardia scabra</i>	A
Rocket, London	<i>Sisymbrium irio</i>	A
Rush skeletonweed ⁴	<i>Chondrilla juncea</i>	B
Saltbush	<i>Atriplex</i> spp.	A
Sheperd's-purse	<i>Capsella bursa-pastoris</i>	A
Spurge, Annual	<i>Euphorbia</i> spp.	A
Stinging nettle ⁴	<i>Urtica dioica</i>	P
Velvetleaf	<i>Abutilon theophrasti</i>	A
Yellow starthistle	<i>Centaurea solstitialis</i>	A

Apply 4-6 pints per acre¹

Arrowwood	<i>Pluchea sericea</i>	A
Canada thistle	<i>Cirsium arvense</i>	P
Giant ragweed	<i>Ambrosia trifida</i>	A
Grey rabbitbrush	<i>Chrysothamnus nauseosus</i>	P
Little mallow	<i>Malva parviflora</i>	B
Milkweed	<i>Asclepias</i> spp.	P
Primrose	<i>Oenothera kunthiana</i>	P
Silverleaf nightshade	<i>Solanum eleagnifolium</i>	P
Sowthistle	<i>Sonchus</i> spp.	A
Texas thistle	<i>Cirsium texanum</i>	P

VINES AND BRAMBLES

Apply 1 pint per acre

Field bindweed	<i>Convolvulus arvensis</i>	P
Hedge bindweed	<i>Calystegia sepium</i>	A

Apply 2-3 pints per acre¹

Wild buckwheat	<i>Polygonum convolvulus</i>	P
----------------	------------------------------	---

Apply 3-4 pints per acre¹

Greenbriar	<i>Smilax</i> spp.	P
Honeysuckle	<i>Lonicera</i> spp.	P
Morningglory	<i>Ipomoea</i> spp.	A/P
Poison ivy	<i>Rhus radicans</i>	P
Redvine	<i>Brunnichia cirrhosa</i>	P
Wild rose	<i>Rosa</i> spp.	P
Including:		P
Multiflora rose	<i>Rosa multiflora</i>	
McCartney rose	<i>Rosa bracteata</i>	P

Apply 4-6 pints per acre¹

*Kudzu ³	<i>Pueraria lobata</i>	P
---------------------	------------------------	---

COMMON NAME	SPECIES	GROWTH HABIT ²
Trumpet creeper	<i>Campsis radicans</i>	P
Virginia creeper	<i>Parthenocissus quinquefolia</i>	P
Wild grape	<i>Vitis</i> spp.	P

BRUSH SPECIES

Apply 4-6 pints per acre ¹		
American beech	<i>Fagus grandifolia</i>	P
Ash	<i>Fraxinus</i> spp.	P
Bald cypress	<i>Taxodium distichum</i>	P
Bigleaf maple	<i>Acer macrophyllum</i>	P
Black locust ⁵	<i>Robinia pseudoacacia</i>	P
Black gum	<i>Nyssa sylvatica</i>	P
Box elder	<i>Acer negundo</i>	P
Cherry	<i>Prunus</i> spp.	P
Chinaberry	<i>Melia azadarach</i>	P
Dogwood	<i>Cornus</i> spp.	P
Elm ⁶	<i>Ulmus</i> spp.	P
Hawthorn	<i>Crataegus</i> spp.	P
Hickory	<i>Carya</i> spp.	P
Honeylocust ⁵	<i>Gleditsia triacanthos</i>	P
Maple	<i>Acer</i> spp.	P
Mulberry	<i>Morus</i> spp.	P
Oak	<i>Quercus</i> spp.	P
Persimmon	<i>Diospyros virginiana</i>	P
*Pine ⁵	<i>Pinus</i> spp.	P
Poplar	<i>Populus</i> spp.	P
Privet	<i>Ligustrum vulgare</i>	P
Red Alder	<i>Alnus rubra</i>	P
Red Maple	<i>Acer rubrum</i>	P
Russian Olive	<i>Eleagnus angustifolia</i>	P
Sassafras	<i>Sassafras albidum</i>	P
Sourwood	<i>Oxydendrum arboreum</i>	P
Sweetgum	<i>Liquidambar styraciflua</i>	P
*Water willow	<i>Justica americana</i>	P
Willow	<i>Salix</i> spp.	P
Yellow poplar	<i>Liriodendron tulipifera</i>	P

¹The higher rates should be used where heavy or well-established infestations occur.

²Growth Habit – A = Annual, B = Biennial, P = Perennial

³Use a minimum of 75 GPA - Control of established stands may require repeat applications.

⁴For best results early postemergence applications are required.

⁵Tank mix with glyphosate or triclopyr.

⁶Tank-mix with glyphosate.

*Not approved for use in California.

ADJUVANTS

For optimal postemergence performance of **Alligare Ecomazapyr 2 SL**, the addition of an adjuvant to the spray solution is essential to aid in the deposition and uptake of the herbicide. For this purpose, **ONLY** use spray adjuvants that are approved or appropriate for aquatic use.

Nonionic Surfactants: Use a nonionic surfactant at 0.25% v/v or higher (depending on surfactant manufacturer's label) of the spray solution (0.25% v/v is equivalent to 1 quart in 100 gallons). For best results, select a nonionic surfactant with HLB (hydrophilic to lipophilic balance) ratio between 12 and 17 with at least 70% surfactant in the formulated product. Alcohols, fatty acids, horticultural spray oils, ethylene glycol or diethylene glycol should not be considered as surfactants to meet these requirements.

Methylated Seed Oils or Vegetable Oil Concentrates: Methylated seed oil or vegetable oil concentrate may be used at 1.5 to 2 pints per acre. When using spray volumes greater than 30 gallons per acre, mix methylated seed oil or vegetable oil concentrate at a rate of 1% of the total spray volume.

Silicone Based Surfactants: Silicone-based surfactants allow greater spreading of the spray droplet on the leaf surface as compared to conventional nonionic surfactants. However, some silicone-based surfactants may dry too quickly and limit herbicide uptake. Refer to the surfactant manufacturer's label for specific recommendations.

Invert emulsions: **Alligare Ecomazapyr 2 SL** can be applied as an invert emulsion (water-in-oil emulsion) designed to minimize spray drift and spray run-off, thereby delivering more herbicide to the target foliage. The spray emulsion may be formed in a single tank (batch mixing) or injected (in-line mixing). Refer to the invert chemical label for proper mixing directions.

Other: Anti-foam agents, spray pattern indicators or drift reduction agents may also be used if necessary or desired. Refer to the adjuvant manufacturer's label for specific recommendations.

AQUATIC USES, PRECAUTIONS AND RESTRICTIONS

Alligare Ecomazapyr 2 SL may only be used for application to control undesirable emergent and floating aquatic vegetation in and around standing and flowing water, including estuarine and marine sites. **Alligare Ecomazapyr 2 SL** may also be used to control undesirable vegetation growing in or around surface water in wetland, riparian and terrestrial habitats where applications may inadvertently contact surface water.

DO NOT apply more than 6 pints of product (1.5 pounds acid equivalent) per acre per year.

Aerial application is restricted to helicopter equipment only.

Alligare Ecomazapyr 2 SL can only be applied by federal or state agencies, such as Water Management District personnel, municipal officials and the U.S. Army Corps of Engineers, or those applicators who are licensed or certified as aquatic pest control applicators and are authorized by the state or local government.

Treatment of other-than-non-native, invasive species is limited to those that have been determined to be nuisance vegetation by a federal or state government entity.

Applications to private waters: **Alligare Ecomazapyr 2 SL** may be applied to private non-flowing waters, such as ponds, lakes and drainage ditches where there is minimal or no outflow to public waters.

Applications to public waters: **Alligare Ecomazapyr 2 SL** may be applied to public waters such as ponds, lakes, reservoirs, marshes, bayous, drainage ditches, canals, streams, rivers and other slow-moving or non-moving waters for control of aquatic weeds or of riparian and wetland weed species.

Consult local state fish and game agency and water control authorities before applying this product to any public waters. Permits may be required to treat such water.

Recreational Use of Water in Treated Areas: There are no restrictions on the use of water in treated areas for recreational purposes, including swimming and fishing.

Livestock Use of Water in/from Treated Areas:

There are no restrictions on livestock consumption of water from treated areas.

Precautions for Potable Water Intakes: DO NOT apply Alligare Ecomazapyr 2 SL directly to waters within one-half mile upstream of an active potable water intake in flowing water (i.e., river, stream, etc.) or within one-half mile of an active potable water intake in a standing body of water such as lake, pond or reservoir. To make aquatic applications around and within one-half mile of active potable water intakes, the water intake must be turned off during application and for at least 48 hours after the treatment. Such aquatic applications may be made only where there are alternative water sources or holding ponds that can be used while active potable water intakes are turned off during the minimum 48 hours post-treatment period. Note: Existing potable water intakes which are no longer in use, such as those replaced by connections to wells or municipal water systems, are not considered to be active potable water intakes. This restriction does not apply to intermittent, inadvertent overspray of water in terrestrial use sites.

APPLICATION TO WATERS USED FOR IRRIGATION

DO NOT use water treated with Alligare Ecomazapyr 2 SL for irrigation purposes for 120 days after treatment or until Alligare Ecomazapyr 2 SL residue levels are determined by laboratory analysis, or other appropriate means of analysis, to be 1.0 ppb or less.

Seasonal Irrigation Waters: Alligare Ecomazapyr 2 SL may be applied during the off-season to surface waters that are used for irrigation on a seasonable basis, provided at least 120 days elapses between Alligare Ecomazapyr 2 SL treatment and the first use of treated water for irrigation or until Alligare Ecomazapyr 2 SL residue levels are determined by laboratory analysis, or other appropriate means of analysis, to be 1.0 ppb or less.

Irrigation Canals/Ditches: DO NOT apply Alligare Ecomazapyr 2 SL to irrigation canals/ditches unless the 120-day restriction on irrigation water usage can be observed or Alligare Ecomazapyr 2 SL residue levels are determined by laboratory analysis, or other appropriate means of analysis, to be 1.0 ppb or less. DO NOT apply Alligare Ecomazapyr 2 SL to dry irrigation canals/ditches.

Quiescent or Slow Moving Waters: In lakes and reservoirs DO NOT apply Alligare Ecomazapyr 2 SL within one (1) mile of an active irrigation water intake during the irrigation season. Applications less than one (1) mile from an inactive irrigation water intake may be made during the off-season, provided the irrigation intake will remain inactive for at least 120 days after treatment or until Alligare Ecomazapyr 2 SL residue levels are determined by laboratory analysis, or other appropriate means of analysis, to be 1.0 ppb or less.

Moving Waters: DO NOT apply Alligare Ecomazapyr 2 SL within one-half mile downstream of an active irrigation water intake. When making applications upstream from an active irrigation water intake, the intake must be turned off for a period of time sufficient to allow the upstream treated water to completely flow past the irrigation intake. Shut off time will be determined by the speed of water flow and the distance and length of water treated upstream from the intake. Consult local, state and/or federal authorities before making any applications upstream from an active irrigation water intake.

TANK MIXES

Alligare Ecomazapyr 2 SL may be tank-mixed with other aquatic-use herbicides to broaden the spectrum of control of emergent and floating aquatic vegetation provided that the tank mix product label does not prohibit such mixing. Consult manufacturer's labels for specific rates and weeds controlled. Always follow the more restrictive label when making an application involving tank-mixes.

AERIAL APPLICATIONS

Exercise all precautions to minimize or eliminate spray drift. Aerial applications may only be made by helicopter; applications may not be made by airplane. Helicopters can be used to apply **Alligare Ecomazapyr 2 SL**; however, **DO NOT** apply by helicopter unless appropriate buffer zones can be maintained to prevent spray drift out of the target area or where damage to desirable vegetation can be tolerated. Helicopters equipped with a Microfoil™ boom, Thru-Valve™ boom or raindrop nozzles must be used and calibrated. Unless applying with a Microfoil boom, use a drift control agent at the recommended label rate. To avoid drift, **DO NOT** make applications during inversion conditions, when winds are gusty, or during any other conditions that promote spray drift. Side trimming is not recommended with **Alligare Ecomazapyr 2 SL** unless death of treated vegetation is acceptable.

Uniformly apply **Alligare Ecomazapyr 2 SL** in 5 to 30 gallons of water per acre. Use a nonionic surfactant, methylated seed oil or silicone-based surfactant (see the **ADJUVANTS** section of this label for specific recommendations). An anti-foam agent may be added, if needed.

Thoroughly clean application equipment, including landing gear, by thoroughly flushing with water immediately after using this product. Prolonged exposure of uncoated/unpainted steel (except stainless steel) surfaces to this product may result in corrosion and failure of the exposed part. Maintaining painted surfaces may prevent corrosion.

GROUND APPLICATIONS

FOLIAR APPLICATIONS:

For Low Volume Foliar Applications: Use equipment calibrated to deliver 5 to 20 gallons of spray solution per acre. Thoroughly mix 0.5 to 5% (v/v) **Alligare Ecomazapyr 2 SL** in water plus surfactant (see the **ADJUVANTS** section of this label for recommendations). Use an anti-foam agent at the recommended rate, if needed. For difficult to control species (see **AQUATIC WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL** section and the **ADDITIONAL WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL** section for relative susceptibility of weed species), apply the higher concentrations of herbicide and/or spray volumes but **DO NOT** apply more than 6 pints of **Alligare Ecomazapyr 2 SL** per acre. Excessive wetting of foliage is not recommended. See the **SPRAY SOLUTION MIXING GUIDE** below for some suggested volumes of **Alligare Ecomazapyr 2 SL** and water.

Select appropriate nozzles to avoid over-application. Proper application is critical to ensure desirable results. Optimum results are achieved when the spray covers the crown and approximately 70 percent of the plant. The use of a flat fan nozzle tip with a spray angle of 40 degrees or less will aid in proper deposition.

Recommended nozzle tip sizes include 4004E or 1504E. For a straight stream and cone pattern, use adjustable cone nozzles such as 5500 X3 or 5500 X4. Attaching a rollover valve onto a Spraying Systems Model 30 gunjet or other similar spray guns allows for the use of both a flat fan and cone tips on the same gun.

Moisten, but do not drench target vegetation. **DO NOT** spray to run off.

For Low Volume Foliar Applications by Backpacks: For low-growing species, spray downward to cover approximately 70% of the plant and the crown.

For target species 4 to 8 feet tall, apply a directed spray in a smooth vertical motion from the crown upward on at least two sides of the target vegetation, making sure to cover the crown whenever possible.

For target species over 8 feet tall, apply a directed spray in a smooth zig-zag motion from the crown upward on at least two sides of the target brush.

For Low Volume Foliar Applications with Hydraulic Handgun Equipment: Use same technique as described above for **Low Volume with Backpacks**.

For broadcast applications, simulate a gentle rain near the top of target vegetation, allowing spray to penetrate the target foliage and contact the crown without run-off onto understory vegetation. **DO NOT** spray to run-off. Herbicide spray that contacts understory vegetation may result in severe injury or death of understory plants.

SPRAY SOLUTION MIXING GUIDE FOR LOW-VOLUME FOLIAR APPLICATIONS

Amount of spray solution	Desired Concentration (fluid volume)				
	0.5%	0.75%	1%	1.5%	5%
(amount of Alligare Ecomazapyr 2 SL to use)					
1 gallon	0.6 oz	0.9 oz	1.3 oz	1.9 oz	6.5 oz
3 gallons	1.9 oz	2.8 oz	3.8 oz	5.8 oz	1.2 pint
4 gallons	2.5 oz	3.8 oz	5.1 oz	7.7 oz	1.6 pint
5 gallons	3.2 oz	4.8 oz	6.5 oz	9.6 oz	2 pints
50 gallons	2 pints	3 pints	4 pints	6 pints	10 pints
100 gallons	4 pints	6 pints	8 pints	6 quarts	5 gallons

2 tablespoons = 1 fluid ounce

For High Volume Foliar Applications: For optimum performance when spraying medium to high-density vegetation, use equipment calibrated to deliver up to 100 gallons of finished spray per acre (GPA). Application volumes exceeding 100 gallons GPA may result in excessive spray run-off, causing injury to desirable ground cover species. Thoroughly mix **Alligare Ecomazapyr 2 SL** in water and include a surfactant (see **ADJUVANTS** section for surfactant recommendations). Use an anti-foam agent according to its label, if needed. For hard-to-control species (see **AQUATIC WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL** section and the **ADDITIONAL WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL** section for relative susceptibility of weeds), use the higher concentrations of the herbicide and/or spray volumes, but **DO NOT** apply more than 6 pints of **Alligare Ecomazapyr 2 SL** per acre. Uniformly cover the foliage of the target vegetation but **DO NOT** apply to run-off.

Side Trimming: **DO NOT** side trim with **Alligare Ecomazapyr 2 SL** unless severe injury or death of the treated vegetation is acceptable. **Alligare Ecomazapyr 2 SL** is readily translocated and can result in the death of the vegetation.

CUT SURFACE TREATMENTS

Alligare Ecomazapyr 2 SL will control undesirable woody vegetation when applied as a water solution to the cambium area of freshly-cut stump surfaces or to cuts on the stem of the target woody vegetation. Applications can be made at any time of the year except during periods of heavy sap flow in the spring. Tree injection and cut stem treatments are most effective in late summer and early fall. **DO NOT** over apply solution causing run-off from the cut surface.

NOTE: Injury may occur to desirable woody plants if the shoots extend from the same root system or their root systems are grafted to those of the treated tree.

CUT SURFACE APPLICATIONS WITH DILUTE AND CONCENTRATE SOLUTIONS:

Mix **Alligare Ecomazapyr 2 SL** as either a concentrated or dilute solution for cut surface treatments. Apply dilute solutions to the cut surface of the stump or to cuts on the stem of the target woody vegetation. Apply concentrate solutions to cuts on the stem. Use of concentrated solutions permits application to fewer cuts on the stem, especially for large diameter trees. Follow the application directions below to determine proper application techniques for each type of solution.

To prepare a dilute solution, mix 8 to 12 fluid ounces of **Alligare Ecomazapyr 2 SL** with one gallon of water. If temperatures are such that freezing of the spray mixture may occur, antifreeze (ethylene glycol) may be added according to manufacturer's label to prevent freezing. The use of a surfactant or penetrating agent may improve herbicide uptake through partially callused cambium tissues.

To prepare a concentrated solution, mix 2 quarts of **Alligare Ecomazapyr 2 SL** with no more than 1 quart water.

For cut stump treatments: Spray or brush the solution onto the cambium area of the freshly cut stump surface. Thoroughly wet the entire cambium area (the wood just inside the bark of the stump).

For cut stem (injection, hack and squirt) treatments:

Dilute Solutions: Using standard injection equipment, apply 1 milliliter of solution at each injection site around the tree with no more than one-inch intervals between cut edges. Insure that the injector completely penetrates the bark at each injection site.

Concentrate Solutions: Using standard injection equipment, apply 1 milliliter of solution at each injection site. Make at least one injection cut for every 3 inches of Diameter at Breast Height (DBH) on the target tree. For example, a 3-inch DBH tree will receive 1 injection cut while a 6-inch DBH tree will receive 2 injection cuts. On trees requiring more than one injection site, place the injection cuts at approximately equal intervals around the tree.

Frill or girdle treatments: Use a hatchet, machete or chainsaw to make cuts through the bark around the tree to expose the cambium. The cuts should angle downward into the cambium sufficiently deep to expose at least two growth rings. Use a spray applicator or brush to apply a 25% to 100% solution of **Alligare Ecomazapyr 2 SL** into each cut until thoroughly wet. Avoid applying so much herbicide that runoff to the ground or water occurs.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: **DO NOT** store below 10°F.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Nonrefillable container (2.5, 30, 265 gallons): Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying.

(Nonrefillable \leq 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

(Nonrefillable $>$ 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Refillable container (15 gallon): Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

IMPORTANT: Read the entire **DIRECTIONS FOR USE** and the **CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY** before using this product. If terms are not acceptable, return the unopened product container at once.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

To the extent consistent with applicable law, upon purchase or use of this product, purchaser and user agree to the following terms:

Warranty: Alligare, LLC (the Company) warrants that this product conforms to the chemical description on the label in all material respects and is reasonably fit for the purpose referred to in the directions for use, subject to the exceptions noted below, which are beyond the Company's control. To the extent consistent with applicable law, the Company makes no other representation or warranty, express or implied, concerning the product, including no implied warranty of merchantability or fitness for a particular purpose. No such warranty shall be implied by law, and no agent or representative is authorized to make any such warranty on the Company's behalf.

Terms of Sale: The Company's directions for use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, and the manner of use or application (including failure to adhere to label directions), all of which are beyond the Company's control. To the extent consistent with applicable law, all such risks are assumed by the user.

Limitation of Liability: To the extent consistent with applicable law, the exclusive remedy against the Company for any cause of action relating to the handling or use of this product is a claim for damages, and in no event shall damages or any other recovery of any kind exceed the price of the product which caused the alleged loss, damage, injury or other claim. To the extent consistent with applicable law, under no circumstances shall the Company be liable for any special, indirect, incidental or consequential damages of any kind, including loss of profits or income, and any such claims are hereby waived. Some states do not allow the exclusion or limitation of incidental or consequential damages.

The Company and the seller offer this product, and the purchaser and user accept this product, subject to the foregoing warranty, terms of sale and limitation of liability, which may be varied or modified only by an agreement in writing signed on behalf of the Company by an authorized representative.

Microfoil is a trademark of Rhone Poulenc Ag. Company.
Thru-Valve is a trademark of Waldrum Specialties.

EPA 20080923



PROTECTING:
RIGHT-OF-WAY
UTILITIES
RAILROAD
RANGE & PASTURELAND
FORESTRY
AQUATIC
INDUSTRIAL

June 3, 2008

To Whom It May Concern:

RE: Letter of Authorization

Dear Sir or Madam:

Please let this letter serve to confirm that Pyxis Regulatory Consulting, Inc. is authorized to act as agent for Alligare, LLC (EPA Company Number 81927), before the U.S. Environmental Protection Agency and state governmental agencies in all matters regarding our pesticide registrations pursuant to the Federal Insecticide, Fungicide and Rodenticide Act ("FIFRA"), 7 U.S.C. § 136 et seq. and state law.

If you have any questions, please do not hesitate to contact me.

Sincerely,

Jay Golz
Vice-President & COO

cc: Pyxis Regulatory Consulting, Inc.



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Mr. Michael Kellogg
Pyxis Regulatory Consulting, Inc
4110 136th N.W
Gig Harbor, WA 98332

SEP 11 2008

Subject: Label Notification(s) for Pesticide Registration Notice 2007-4

Dear Mr. Kellogg:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 dated August 25, 08 for:

EPA Registration 81927-22 Alligare Ecomazapyr 2SL

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-4 and finds that the label change(s) requested falls within the scope of PRN-2007-4. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact me directly at 703-305-6249 or Banza Djapao of my staff at 703-305-7269.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

NOTIFICATION**Alligare Ecomazapyr 2 SL**

SEP 11 2008

Alligare Ecomazapyr 2 SL may only be used for application to control undesirable emergent and floating aquatic vegetation in and around standing and flowing water, including estuarine and marine sites. **Alligare Ecomazapyr 2 SL** may also be used to control undesirable vegetation growing in or around surface water in wetland, riparian and terrestrial habitats where applications may inadvertently contact surface water. **Alligare Ecomazapyr 2 SL** may also be used for cut stump, cut stem and frill and girdle treatments within aquatic sites.

ACTIVE INGREDIENT:

Isopropylamine salt of Imazapyr (2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid)*27.8%

OTHER INGREDIENTS72.2%

TOTAL100.0%

*Equivalent to 22.6% 2-[4,5-dihydro-4-methyl-4-(1-methylethyl)-5-oxo-1H-imidazol-2-yl]-3-pyridinecarboxylic acid or 2 pounds acid equivalent per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION/PRECAUCIÓN**

PRECAUCION AL USUARIO: Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• DO NOT induce vomiting unless told to so by the poison control center or doctor.• DO NOT give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For medical emergencies involving this product, call 1-800-424-9300.	


EPA Reg. No. 81927-22

EPA Est. No.

Manufactured For:

Alligare, LLC
13 N. 8th Street
Opelika, AL 36801

Net Contents:



PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION! Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing and wash before reuse.

Personal Protective Equipment (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistant category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material.
- Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions are given for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS

DO NOT apply to water except as specified in this label. Treatment of aquatic weeds may result in oxygen depletion or loss due to decomposition of dead plants. This oxygen loss may cause the suffocation of some aquatic organisms. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

This pesticide is toxic to vascular plants and should be used strictly in accordance with the drift precautions on the label.

PHYSICAL AND CHEMICAL HAZARDS

Spray solutions of **Alligare Ecomazapyr 2 SL** should be mixed, stored and applied only in stainless steel, fiberglass, plastic and plastic-lined steel containers.

DO NOT mix, store or apply **Alligare Ecomazapyr 2 SL** or spray solutions of **Alligare Ecomazapyr 2 SL** in unlined steel (except stainless steel) containers or spray tanks.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Keep containers closed to avoid spills and contamination.

GENERAL INFORMATION

Alligare Ecomazapyr 2 SL is an aqueous solution intended to be mixed with water and surfactant(s) for application to control floating and emergent undesirable vegetation (see **AQUATIC WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL** section and the **ADDITIONAL WEEDS**

CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL section) in or near bodies of water which may be flowing, non-flowing or transient. **Alligare Ecomazapyr 2 SL** may be applied to aquatic sites that include lakes, rivers, streams, ponds, seeps, drainage ditches, canals, reservoirs, swamps, bogs, marshes, estuaries, bays, brackish water, transitional areas between terrestrial and aquatic sites and seasonal wet areas. See **AQUATIC USES, PRECAUTIONS AND RESTRICTIONS** section of this label for precautions, restrictions, and instructions on aquatic uses. **Alligare Ecomazapyr 2 SL** may also be used for cut stump, cut stem and frill and girdle treatments within aquatic sites (see **AERIAL APPLICATIONS AND GROUND APPLICATIONS** sections for additional details).

DO NOT treat more than one half of the surface area of the water in a single operation and wait at least 10 to 14 days between treatments. Begin treatment along the shore and proceed outward in bands to allow aquatic organisms to move into untreated areas.

Read and observe the following directions if aquatic sites are present in terrestrial non-crop areas and are part of the intended treatment area:

Alligare Ecomazapyr 2 SL controls most annual and perennial grasses, broadleaf weeds and many brush and vine species. **Alligare Ecomazapyr 2 SL** will provide some residual control of undesirable species that germinate above the waterline. **Alligare Ecomazapyr 2 SL** controls vegetation by absorption through emergent leaves and stems, from which it is translocated throughout the plant, where it accumulates in rapidly-growing meristematic tissue. Treated plants stop growing soon after treatment. Chlorosis (yellowing of plant tissue) first appears in the newest leaves, and necrosis spreads from this point. In perennials, **Alligare Ecomazapyr 2 SL** is translocated into and kills underground or submerged storage organs to prevent regrowth. Chlorosis and tissue necrosis may not be apparent in some plant species until two or more weeks after application. Complete kill of plants may not occur for several weeks. Applications of **Alligare Ecomazapyr 2 SL** are rainfast within one hour after treatment.

Alligare Ecomazapyr 2 SL will not control plants which are completely submerged or have a majority or their foliage under water.

Alligare Ecomazapyr 2 SL must be applied to the emergent foliage of the target vegetation, since it has little-to-no activity on submerged vegetation. Residual concentrations of **Alligare Ecomazapyr 2 SL** from direct application to water are not expected to be sufficient to provide control of undesirable vegetation. Apply **Alligare Ecomazapyr 2 SL** so as to maximize spray contact onto the target vegetation while minimizing overspray onto adjacent or underlying water. For optimum performance, include a surfactant (see **ADJUVANTS** section for specific recommendations) in the spray solution and apply to actively growing weeds. **Alligare Ecomazapyr 2 SL** may be applied in spot treatments by using low-volume directed application techniques or broadcast by using ground equipment, watercraft or helicopter.

Apply Alligare Ecomazapyr 2 SL with surface or helicopter application equipment in at least 5 gallons of water per acre. When applying by helicopter, follow directions under the **AERIAL APPLICATIONS** section of this label, otherwise refer to section on **GROUND APPLICATIONS** when using surface equipment.

Applications to moving bodies of water should be made while traveling upstream to prevent concentration of this herbicide in water. **DO NOT** apply to bodies of water or portions of bodies of water where emergent and/or floating weeds do not exist.

When target vegetation covers a large percentage of the surface area of impounded water, apply **Alligare Ecomazapyr 2 SL** to alternating strips (i.e., **DO NOT** apply to every other swath) to avoid oxygen depletion due to decaying vegetation, since oxygen depletion may result in the suffocation of sensitive aquatic organisms. **DO NOT** treat more than one half of the surface area of the water in a single operation and wait at least 10 to 14 days before retreatment. Begin treatment along the shore and proceed outward in alternating swaths as described above to allow aquatic organisms to move into untreated areas.

Avoid wash-off of sprayed foliage by spray boat or recreational boat backwash for one hour after application.

Apply **Alligare Ecomazapyr 2 SL** at 2 to 6 pints per acre, depending on weed species and weed density. Use higher rates for heavy weed pressure. Refer to the **AQUATIC WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL** section and the **ADDITIONAL WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL** section of this label for specifics. **DO NOT** exceed 6 pints per acre (1.5 lb ai/A) per year.

Alligare Ecomazapyr 2 SL may also be applied as a draw down treatment in areas described above. Apply **Alligare Ecomazapyr 2 SL** to weeds after water has been drained and allow 14 days before reintroduction of water.

PRECAUTIONS FOR AVOIDING INJURY TO NON-TARGET PLANTS

Alligare Ecomazapyr 2 SL can occasionally affect non-target or desirable vegetation by root uptake of the herbicide from treated soil. Injury or loss of non-target plants may result if **Alligare Ecomazapyr 2 SL** is applied onto or near desirable plants, or to areas where their roots extend, or in areas where treated soil may be washed or moved into contact with their root zone.

When making applications along shorelines where desirable plants may be present, exercise caution to avoid spray contact onto their foliage or spray application to the soil in which they are rooted. Shoreline plants with roots that extend into the waters where **Alligare Ecomazapyr 2 SL** has been applied will generally not be adversely affected.

If treated vegetation is to be removed from the application site, **DO NOT** use the vegetative matter as mulch or compost on or around desirable species.

IMPORTANT

DO NOT use on food crops. **DO NOT** apply this product within one-half mile upstream of an active potable water intake in flowing water (i.e., river, stream, etc.) or within one-half mile of an active potable water intake in a standing body of water, such as a lake, pond or reservoir. **DO NOT** apply to water used for irrigation except as described in **APPLICATION TO WATERS USED FOR IRRIGATION** section of this label. Keep away from fertilizers, insecticides, fungicides, and seeds. **DO NOT** drain or flush equipment on or near desirable plants, or onto areas where their roots may extend, or in locations where the chemical or treated soil may be washed or moved into contact with their roots. **DO NOT** use on lawns, walks, driveways, tennis courts or similar areas where roots of desirable vegetation may extend and be exposed to potential injury and/or mortality from root uptake of **Alligare Ecomazapyr 2 SL**. **DO NOT** side trim desirable vegetation with this product unless severe injury or plant death is acceptable. Prevent drift of spray to desirable plants.

Clean application equipment after using this product by thoroughly flushing with water.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are **NOT** within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Non-crop weed control is not within the scope of the Worker Protection Standard. See the **GENERAL INFORMATION** section of this label for a description of non-crop sites.

DO NOT enter treated areas without protective clothing until sprays have dried.

SPRAY DRIFT MANAGEMENT

Spray Drift: Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determines the potential for spray drift. The applicator and the entity authorizing spraying are responsible for considering all these factors when making decisions.

Spray drift from applying this product may result in damage to sensitive plants adjacent to the treatment area. Only apply this product when the potential for drift to these and other adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, or non-target crops) is minimal. **DO NOT** apply when the following conditions exist that increase the likelihood of spray drift from intended targets: high or gusty winds, high temperatures, low humidity, and temperature inversions.

To minimize spray drift, the applicator should be familiar with and take into account the following drift reduction advisory information. Additional information may be available from state enforcement agencies or the Cooperative Extension on the application of this product.

The best drift management strategy and most effective way to reduce drift potential are to apply large droplets that provide sufficient coverage and control. Applying larger droplets reduces the drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (**see WIND, TEMPERATURE AND HUMIDITY, AND TEMPERATURE INVERSIONS**).

Controlling Droplet Size:

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - **DO NOT** exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- Number of Nozzles - Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is recommended practice. Significant deflection from the horizontal will reduce droplet size and increase drift potential.
- Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift. **DO NOT** use nozzles producing a mist droplet spray.

Application Height: Making applications at the lowest possible height (helicopter, ground driven spray bloom) that is safe and practical reduces exposure of droplets to evaporation and wind.

Swath Adjustment: When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the treatment area, the applicator must compensate for this displacement by adjusting the path of the application equipment (e.g. aircraft, ground) upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller droplets, etc.).

Wind: Drift potential is lowest between wind speeds of 3-10 mph. However, many factors, including droplet size and equipment type, determine drift potential at any given speed. Application should be avoided below 3 mph due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions: Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud, which can move in unpredictable directions due to the light variable winds common during temperature inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Wind Erosion: Avoid treating powdery dry or light sandy soils when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation.

Aerial Application Methods and Equipment (Helicopters ONLY): Water Volume: Use 2 or more gallons of water per acre. The actual minimum spray volume per acre is determined by the spray equipment used. Use adequate spray volume to produce accurate and uniform distribution of spray particles over the treated area and to avoid spray drift.

Managing Spray Drift from Aerial Applications: Applicators must follow these requirements to avoid off-target drift movement: 1) boom length- the distance of the outermost nozzles on the boom must not exceed $\frac{1}{4}$ the length of the rotor, 2) nozzle orientation- nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees, and 3) application height- without compromising helicopter safety, applications should be made at a height of 10 feet or less above the crop canopy or tallest plants. Applicators must follow the most restrictive use cautions to avoid drift hazards, including those found in this labeling as well as applicable state and local regulations and ordinances.

Ground Application (Broadcast): Water Volume: Use 5 or more gallons of water per acre. The actual minimum spray volume per acre is determined by the spray equipment used. Use adequate spray volume to provide accurate and uniform distribution of spray particles over the treated area and to avoid spray drift.

AQUATIC SPECIES CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL

Alligare Ecomazapyr 2 SL will control the following target species when used as specified in the table below. Rate instructions are expressed as product volume for broadcast applications and as a % solution for directed applications including spot treatments. For % solution applications, **DO NOT** apply more than the equivalent of 3 quarts of Alligare Ecomazapyr 2 SL per acre.

COMMON NAME	SCIENTIFIC NAME	INSTRUCTIONS
Floating Species		
*Duckweed	<i>Lemna minor</i>	2-3 pints/acre (1% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
*Duckweed, Giant	<i>Spirodela polyrriza</i>	2-3 pints/acre (1% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
*Frogbit	<i>Limnobium spongia</i>	1-2 pints/acre (0.5% solution) in 100 GPA water.

COMMON NAME	SCIENTIFIC NAME	INSTRUCTIONS
*Spatterdock	<i>Nuphar luteum</i>	Completely cover all actively-growing emergent foliage. Apply a tank-mix of 2-4 pints/acre Alligare Ecomazapyr 2 SL plus 4-6 pints/acre glyphosate (0.5% Alligare Ecomazapyr 2 SL plus 1.5% glyphosate) in 100 GPA water for best control. Completely cover all actively-growing emergent foliage.
*Water Hyacinth	<i>Eichhornia crassipes</i>	1-2 pints/acre (0.5% solution) in 100 GPA water to actively growing foliage.
*Water Lettuce	<i>Pistia stratiotes</i>	1-2 pints/acre (0.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.

Emerged Species

*Alligatorweed	<i>Alternanthera philoxeroides</i>	1-4 pints/acre (0.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage. DO NOT apply as a tank-mix with glyphosate, since doing so may reduce control.
*Arrowhead, Duck-potato	<i>Sagittaria</i> spp.	1-2 pints/acre (0.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
*Bacopa, lemon	<i>Bacopa</i> spp.	1-2 pints/acre (0.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
*Parrot feather	<i>Myriophyllum aquaticum</i>	Apply 2-4 pints in 100 GPA water to actively growing emergent foliage. Foliage must be above water for sufficient uptake.
*Pennywort	<i>Hydrocotyle</i> spp.	1-2 pints/acre (0.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
*Pickerelweed	<i>Pontederia cordata</i>	2-3 pints/acre (1% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
*Taro, wild; Dasheen; Elephant's Ear; Coco Yam	<i>Colocasia esculentum</i>	4-6 pints/acre (1.5% solution) in 100 GPA with a high quality 'sticker' adjuvant. Completely cover all actively-growing emergent foliage.
*Water lily	<i>Nymphaea odorata</i>	2-3 pints/acre (1% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage.
*Water Primrose	<i>Ludwigia uruguayensis</i>	4-6 pints/acre (1.5% solution) in 100 GPA water. Completely cover all actively-growing emergent foliage. DO NOT apply as a tank-mix with glyphosate, since doing so may reduce control.

Terrestrial/Marginal

*Soda Apple, aquatic; Nightshade	<i>Solanum tampicense</i>	2 pints/acre applied to foliage.
*Bamboo, Japanese	<i>Phyllostachys</i> spp.	3-4 pints/acre applied to actively-growing foliage before plants set seed heads. The greater the amount of foliage present at treatment, the greater the exposure to the herbicide for plant uptake, which will result in greater root kill.
Brazilian Pepper;	<i>Schinus terebinthifolius</i>	2-4 pints/acre applied to foliage.

COMMON NAME	SCIENTIFIC NAME	INSTRUCTIONS
Christmasberry Cattail	<i>Typha</i> spp.	2-4 pints (1% solution) applied to actively-growing, green foliage after full leaf elongation. Lower rates will control cattail in the north; higher rates are needed in the south.
Chinese Tallow Tree Cogon grass	<i>Sapium sebiferum</i> <i>Imperata cylindrica</i>	16-24 oz applied to foliage. Burn foliage, till area, then treat in fall at 2 quarts/acre Alligare Ecomazapyr 2 SL plus MSO applied to new growth.
Cordgrass, prairie *Cutgrass *Elephant Grass; Napier Grass *Flowering rush Giant Reed, Wild Cane	<i>Spartina</i> spp. <i>Zizaniopsis miliacea</i> <i>Pennisetum purpureum</i> <i>Butumu typla</i> <i>Arundo donax</i>	4-6 pints applied to actively growing foliage. 4-6 pints applied to actively growing foliage. 3 pints/acre applied to actively growing foliage.
*Golden Bamboo	<i>Phyllostachys aurea</i>	2-3 pints applied to actively growing foliage. 4-6 pints/acre applied in spring to actively growing foliage.
Junglerice Knapweeds	<i>Echinolchloa colonum</i> <i>Centaurea species</i>	3-4 pints/acre applied to actively-growing foliage before plants set seed heads. The greater the amount of foliage present at treatment, the greater the exposure to the herbicide for plant uptake, which will result in greater root kill. 3-4 pints applied to actively growing foliage. Russian Knapweed- Apply 2-3 pints plus 1 quart/acre MSO in the fall after senescence begins.
Knotweed, Japanese (see Fallopia japonica) Melaleuca; Paperbark Tree	<i>Polygonum cuspidatum</i> <i>Melaleuca quinquenervia</i>	3-4 pints/acre applied postemergence to actively growing foliage. For established stands, apply 6 pints/acre Alligare Ecomazapyr 2 SL plus 6 pints/acre glyphosate plus adjuvant. For best results, use 4 quarts/A methylated seed oil for adjuvant. For ground foliar application, uniformly apply to completely cover all actively-growing foliage. For aerial foliar application, apply in at least two passes in a cross pattern at 10 GPA. For spot treatment, apply a 25% solution of Alligare Ecomazapyr 2 SL plus 25% solution of glyphosate plus 1.25% MSO in water as a frill or stump treatment.
*Nutgrass; Kili'p'opu	<i>Cyperus rotundus</i>	2 pints Alligare Ecomazapyr 2 SL plus 1 quart/acre MSO applied early postemergence.
*Nutsedge	<i>Cyperus</i> spp.	2-3 pints postemergence to foliage or pre-emergence incorporated. Non-incorporated preemergence applications will not provide control.
Phragmites; Common Reed	<i>Phragmites australis</i>	4-6 pints/acre applied to actively-growing, green foliage after leaf elongation. Completely cover all actively-growing foliage. If stand has a substantial amount of old stem tissue, mow or burn, allow to regrow to approximately 5' tall before treatment. Lower rates will control phragmites in the north; higher rates are needed in the south.
*Poison Hemlock	<i>Conium maculatum</i>	2 pints Alligare Ecomazapyr 2 SL plus 1 quart/acre MSO applied preemergence or early

COMMON NAME	SCIENTIFIC NAME	INSTRUCTIONS
Purple Loosestrife	<i>Lythrum salicaria</i>	postemergence up to rosette stage before flowering.
Reed canarygrass	<i>Phalaris arundinacea</i>	1 pint/acre applied to actively growing foliage.
Rose, swamp	<i>Rosa palustris</i>	3-4 pints/acre applied to actively growing foliage.
Russian-Olive	<i>Elaeagnus angustifolia</i>	2-3 pints/ acre applied to actively growing foliage.
Saltcedar; Tamarisk	<i>Tamarix species</i>	2-4 pints/acre or 1% solution applied to foliage. Aerially apply 2 quarts Alligare Ecomazapyr 2 SL plus 0.25% v/v NIS to actively growing foliage during flowering. For spot spraying, use a 1% solution of Alligare Ecomazapyr 2 SL plus 0.25% v/v NIS and spray to wet foliage. After application wait at least two years before disturbing treated saltcedar. Earlier disturbance can reduce overall control.
Smartweed	<i>Polygonum spp.</i>	2 pints/acre applied early postemergence.
Sumac	<i>Rhus spp.</i>	2-3 pints/acre applied to foliage.
Swamp Morning Glory;	<i>Ipomoea aquatica</i>	1-2 pints/acre Alligare Ecomazapyr 2 SL plus 1 quart/acre MSO applied at early postemergence.
Water Spinach;		
Kangkong		
Torpedo grass	<i>Panicum repens</i>	4 pints/acre (1-1.5% solution). Completely cover all actively growing foliage.
*White Top; Hoary	<i>Cardaria draba</i>	1-2 pints/acre applied in spring to foliage during flowering.
Cress		
Willow	<i>Salix spp.</i>	2-3 pints/acre Alligare Ecomazapyr 2 SL . Completely cover all actively growing foliage.

* Not approved for use in California

ADDITIONAL WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL

When used as recommended on terrestrial sites, **Alligare Ecomazapyr 2 SL** will provide preemergence or postemergence control with residual control of the following vegetation species. Annual weeds may be controlled by preemergence or postemergence applications of **Alligare Ecomazapyr 2 SL**. For established biennials and perennial vegetation control, postemergence treatments of **Alligare Ecomazapyr 2 SL** are recommended.

The tables below show broadcast rates and indicate relative weed sensitivity. It is important to consider relative weed sensitivity when preparing low volume spray solutions (see **LOW VOLUME** section of **GROUND APPLICATIONS** section), since low volume treatments apply less **Alligare Ecomazapyr 2 SL** per acre than is shown for the broadcast treatments.

The relative sensitivity of the species listed below can also be used to determine the relative risk of causing non-target plant injury if any of those species are considered to be desirable within the treatment area.

Resistant Biotypes: Some weeds listed below may have naturally-occurring biotypes (plants within a given species that have a slightly different but distinct genetic makeup from other plants of that species) that are not effectively controlled by this and/or other herbicides (Oust®) with the ALS/AHAS enzyme-inhibiting mode of action. If naturally-occurring ALS/AHAS-resistant biotypes are present in area, **Alligare Ecomazapyr 2 SL** should be tank-mixed or applied sequentially with a registered herbicide that depends on a different mode of action to ensure control.

GRASSES

COMMON NAME	SPECIES	GROWTH HABIT ²
Apply 2-3 pints per acre¹		
Annual bluegrass	<i>Poa annua</i>	A
Broadleaf signalgrass	<i>Brachiaria platyphylla</i>	A
Canada bluegrass	<i>Poa compressa</i>	P
Downy brome	<i>Bromus tectorum</i>	A
Fescue	<i>Festuca</i> spp.	A/P
Foxtail	<i>Setaria</i> spp.	A
Italian ryegrass	<i>Lolium multiflorum</i>	A
Johnsongrass	<i>Sorghum halepense</i>	P
Kentucky bluegrass	<i>Poa pratensis</i>	P
Lovegrass	<i>Eragrostis</i> spp.	A/P
*Napier grass	<i>Pennisetum purpureum</i>	P
Orchardgrass	<i>Dactylis glomerata</i>	P
Paragrass	<i>Brachiaria mutica</i>	P
Quackgrass	<i>Agropyron repens</i>	P
Sandbur	<i>Cenchrus</i> spp.	A
Sand dropseed	<i>Sporobolus cryptandrus</i>	P
Smooth brome	<i>Bromus inermis</i>	P
Vaseygrass	<i>Paspalum urvillei</i>	P
Wild Oats	<i>Avena fatua</i>	A
Witchgrass	<i>Panicum capillare</i>	A
Apply 3-4 pints per acre¹		
Barnyardgrass	<i>Echinochloa crus-gali</i>	A
Beardgrass	<i>Andropogon</i> spp.	P
Bluegrass, Annual	<i>Poa annua</i>	A
*Bulrush	<i>Scirpus validus</i>	P
Cheat	<i>Bromus secalinus</i>	A
Crabgrass	<i>Digitaria</i> spp.	A
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	A
Fall Panicum	<i>Panicum dichotomiflorum</i>	A
Goosegrass	<i>Eleusine indica</i>	A
Itchgrass	<i>Rottboellia exaltata</i>	A
Lovegrass	<i>Eragrostis</i> spp.	A
*Maidencane	<i>Panicum hemitomon</i>	A
Panicum, Browntop	<i>Panicum fasciculatum</i>	A
Panicum, Texas	<i>Panicum texanum</i>	A
Prairie threeawn	<i>Aristida oligantha</i>	P
Sandbur, Field	<i>Cenchrus incertus</i>	A
Signalgrass	<i>Brachiaria platyphylla</i>	A
Wild barley	<i>Hordeum</i> spp.	A
Wooly Cupgrass	<i>Eriochloa villosa</i>	A
Apply 4-6 pints per acre¹		
Bahiagrass	<i>Paspalum notatum</i>	P
Bermudagrass ³	<i>Cynodon dactylon</i>	P
Big bluestem	<i>Andropogon gerardii</i>	P
Dallisgrass	<i>Paspalum dilatatum</i>	P
Feathertop	<i>Pennisetum villosum</i>	P
Guineagrass	<i>Panicum maximum</i>	P
Saltgrass ³	<i>Distichlis stricta</i>	P
Sand dropseed	<i>Sporobolus cryptandrus</i>	P

COMMON NAME	SPECIES	GROWTH HABIT ²
Sprangletop	<i>Leptochloa</i> spp.	A
Timothy	<i>Phleum pretense</i>	P
Wirestem muhly	<i>Muhlenbergia frondosa</i>	P

BROADLEAF WEEDS

Apply 2-3 pints per acre ¹		
Burdock	<i>Arctium</i> spp.	B
Carpetweed	<i>Mollugo verticillata</i>	A
Carolina geranium	<i>Geranium carolinianum</i>	A
Clover	<i>Trifolium</i> spp.	A/P
Common chickweed	<i>Stellaria media</i>	A
Common ragweed	<i>Ambrosia artemisiifolia</i>	A
Dandelion	<i>Taraxacum officinale</i>	P
Dog fennel	<i>Eupatorium capillifolium</i>	A
Filaree	<i>Erodium</i> spp.	A
Fleabane	<i>Erigeron</i> spp.	A
Hoary vervain	<i>Verbena stricta</i>	P
Indian mustard	<i>Brassica juncea</i>	A
Kochia	<i>Kochia scoparia</i>	A
Lambsquarters	<i>Chenopodium album</i>	A
*Lespedeza	<i>Lespedeza</i> spp.	P
Miners lettuce	<i>Montia perfoliata</i>	A
Mullein	<i>Verbascum</i> spp.	B
Nettleleaf goosefoot	<i>Chenopodium murale</i>	A
Oxeye daisy	<i>Chrysanthemum leucanthemum</i>	P
Pepperweed	<i>Lepidium</i> spp.	A
Pigweed	<i>Amaranthus</i> spp.	A
Puncturevine	<i>Tribulus terrestris</i>	A
Russian thistle	<i>Salsola kali</i>	A
Smartweed	<i>Polygonum</i> spp.	A/P
Sorrell	<i>Rumex</i> spp.	P
Sunflower	<i>Helianthus</i> spp.	A
Sweet clover	<i>Melilotus</i> spp.	A/B
Tansymustard	<i>Descurainia pinnata</i>	A
Western ragweed	<i>Ambrosia psilostachya</i>	P
Wild carrot	<i>Daucus carota</i>	B
Wild lettuce	<i>Lactuca</i> spp.	A/B
Wild parsnip	<i>Pastinaca sativa</i>	B
Wild turnip	<i>Brassica campestris</i>	B
Woollyleaf bursage	<i>Franseria tomentosa</i>	P
Yellow woodsorrel	<i>Oxalis stricta</i>	P

Apply 3-4 pints per acre ¹		
Broom snakeweed ⁴	<i>Gutierrezia sarothrae</i>	P
Bull thistle	<i>Cirsium vulgare</i>	B
Burclover	<i>Medicago</i> spp.	A
Chickweed, Mouseear	<i>Cerastium vulgatum</i>	A
Clover, Hop	<i>Trifolium procumbens</i>	A
Cocklebur	<i>Xanthium strumarium</i>	A
Cudweed	<i>Gnaphalium</i> spp.	A

COMMON NAME	SPECIES	GROWTH HABIT ²
Desert Camelthorn	<i>Alhagi pseudalhagi</i>	P
Dock	<i>Rumex</i> spp.	P
Fiddleneck	<i>Amsinckia intermedia</i>	A
Goldenrod	<i>Solidago</i> spp.	P
Henbit	<i>Lamium alexicaule</i>	A
Knotweed, prostrate	<i>Polygonum aviculare</i>	A/P
Pokeweed	<i>Phytolacca americana</i>	P
Purslane	<i>Portulaca</i> spp.	A
Pusley, Florida	<i>Richardia scabra</i>	A
Rocket, London	<i>Sisymbrium irio</i>	A
Rush skeletonweed ⁴	<i>Chondrilla juncea</i>	B
Saltbush	<i>Atriplex</i> spp.	A
Shepherd's-purse	<i>Capsella bursa-pastoris</i>	A
Spurge, Annual	<i>Euphorbia</i> spp.	A
Stinging nettle ⁴	<i>Urtica dioica</i>	P
Velvetleaf	<i>Abutilon theophrasti</i>	A
Yellow starthistle	<i>Centaurea solstitialis</i>	A

Apply 4-6 pints per acre¹

Arrowwood	<i>Pluchea sericea</i>	A
Canada thistle	<i>Cirsium arvense</i>	P
Giant ragweed	<i>Ambrosia trifida</i>	A
Grey rabbitbrush	<i>Chrysothamnus nauseosus</i>	P
Little mallow	<i>Malva parviflora</i>	B
Milkweed	<i>Asclepias</i> spp.	P
Primrose	<i>Oenothera kunthiana</i>	P
Silverleaf nightshade	<i>Solanum eleagnifolium</i>	P
Sowthistle	<i>Sonchus</i> spp.	A
Texas thistle	<i>Cirsium texanum</i>	P

VINES AND BRAMBLES

Apply 1 pint per acre

Field bindweed	<i>Convolvulus arvensis</i>	P
Hedge bindweed	<i>Calystegia sepium</i>	A

Apply 2-3 pints per acre¹

Wild buckwheat	<i>Polygonum convolvulus</i>	P
----------------	------------------------------	---

Apply 3-4 pints per acre¹

Greenbriar	<i>Smilax</i> spp.	P
Honeysuckle	<i>Lonicera</i> spp.	P
Morningglory	<i>Ipomoea</i> spp.	A/P
Poison ivy	<i>Rhus radicans</i>	P
Redvine	<i>Brunnichia cirrhosa</i>	P
Wild rose	<i>Rosa</i> spp.	P
Including:		P
Multiflora rose	<i>Rosa multiflora</i>	
McCartney rose	<i>Rosa bracteata</i>	P

Apply 4-6 pints per acre¹

*Kudzu ³	<i>Pueraria lobata</i>	P
---------------------	------------------------	---

COMMON NAME	SPECIES	GROWTH HABIT ²
Trumpet creeper	<i>Campsis radicans</i>	P
Virginia creeper	<i>Parthenocissus quinquefolia</i>	P
Wild grape	<i>Vitis</i> spp.	P

BRUSH SPECIES

Apply 4-6 pints per acre ¹		
American beech	<i>Fagus grandifolia</i>	P
Ash	<i>Fraxinus</i> spp.	P
Bald cypress	<i>Taxodium distichum</i>	P
Bigleaf maple	<i>Acer macrophyllum</i>	P
Black locust ⁵	<i>Robinia pseudoacacia</i>	P
Black gum	<i>Nyssa sylvatica</i>	P
Box elder	<i>Acer negundo</i>	P
Cherry	<i>Prunus</i> spp.	P
Chinaberry	<i>Melia azadarach</i>	P
Dogwood	<i>Cornus</i> spp.	P
Elm ⁶	<i>Ulmus</i> spp.	P
Hawthorn	<i>Crataegus</i> spp.	P
Hickory	<i>Carya</i> spp.	P
Honeylocust ⁵	<i>Gleditsia triacanthos</i>	P
Maple	<i>Acer</i> spp.	P
Mulberry	<i>Morus</i> spp.	P
Oak	<i>Quercus</i> spp.	P
Persimmon	<i>Diospyros virginiana</i>	P
*Pine ⁵	<i>Pinus</i> spp.	P
Poplar	<i>Populus</i> spp.	P
Privet	<i>Ligustrum vulgare</i>	P
Red Alder	<i>Alnus rubra</i>	P
Red Maple	<i>Acer rubrum</i>	P
Russian Olive	<i>Eleagnus angustifolia</i>	P
Sassafras	<i>Sassafras albidum</i>	P
Sourwood	<i>Oxydendrum arboreum</i>	P
Sweetgum	<i>Liquidambar styraciflua</i>	P
*Water willow	<i>Justica americana</i>	P
Willow	<i>Salix</i> spp.	P
Yellow poplar	<i>Liriodendron tulipifera</i>	P

¹The higher rates should be used where heavy or well-established infestations occur.

²Growth Habit – A = Annual, B = Biennial, P = Perennial

³Use a minimum of 75 GPA - Control of established stands may require repeat applications.

⁴For best results early postemergence applications are required.

⁵Tank mix with glyphosate or triclopyr.

⁶Tank-mix with glyphosate.

*Not approved for use in California.

ADJUVANTS

For optimal postemergence performance of **Alligare Ecomazapyr 2 SL**, the addition of an adjuvant to the spray solution is essential to aid in the deposition and uptake of the herbicide. For this purpose, **ONLY** use spray adjuvants that are approved or appropriate for aquatic use.

Nonionic Surfactants: Use a nonionic surfactant at 0.25% v/v or higher (depending on surfactant manufacturer's label) of the spray solution (0.25% v/v is equivalent to 1 quart in 100 gallons). For best results, select a nonionic surfactant with HLB (hydrophilic to lipophilic balance) ratio between 12 and 17 with at least 70% surfactant in the formulated product. Alcohols, fatty acids, horticultural spray oils, ethylene glycol or diethylene glycol should not be considered as surfactants to meet these requirements.

Methylated Seed Oils or Vegetable Oil Concentrates: Methylated seed oil or vegetable oil concentrate may be used at 1.5 to 2 pints per acre. When using spray volumes greater than 30 gallons per acre, mix methylated seed oil or vegetable oil concentrate at a rate of 1% of the total spray volume.

Silicone Based Surfactants: Silicone-based surfactants allow greater spreading of the spray droplet on the leaf surface as compared to conventional nonionic surfactants. However, some silicone-based surfactants may dry too quickly and limit herbicide uptake. Refer to the surfactant manufacturer's label for specific recommendations.

Invert emulsions: **Alligare Ecomazapyr 2 SL** can be applied as an invert emulsion (water-in-oil emulsion) designed to minimize spray drift and spray run-off, thereby delivering more herbicide to the target foliage. The spray emulsion may be formed in a single tank (batch mixing) or injected (in-line mixing). Refer to the invert chemical label for proper mixing directions.

Other: Anti-foam agents, spray pattern indicators or drift reduction agents may also be used if necessary or desired. Refer to the adjuvant manufacturer's label for specific recommendations.

AQUATIC USES, PRECAUTIONS AND RESTRICTIONS

Alligare Ecomazapyr 2 SL may only be used for application to control undesirable emergent and floating aquatic vegetation in and around standing and flowing water, including estuarine and marine sites. **Alligare Ecomazapyr 2 SL** may also be used to control undesirable vegetation growing in or around surface water in wetland, riparian and terrestrial habitats where applications may inadvertently contact surface water.

DO NOT apply more than 6 pints of product (1.5 pounds acid equivalent) per acre per year.

Aerial application is restricted to helicopter equipment only.

Alligare Ecomazapyr 2 SL can only be applied by federal or state agencies, such as Water Management District personnel, municipal officials and the U.S. Army Corps of Engineers, or those applicators who are licensed or certified as aquatic pest control applicators and are authorized by the state or local government.

Treatment of other-than-non-native, invasive species is limited to those that have been determined to be nuisance vegetation by a federal or state government entity.

Applications to private waters: **Alligare Ecomazapyr 2 SL** may be applied to private non-flowing waters, such as ponds, lakes and drainage ditches where there is minimal or no outflow to public waters.

Applications to public waters: **Alligare Ecomazapyr 2 SL** may be applied to public waters such as ponds, lakes, reservoirs, marshes, bayous, drainage ditches, canals, streams, rivers and other slow-moving or non-moving waters for control of aquatic weeds or of riparian and wetland weed species.

Consult local state fish and game agency and water control authorities before applying this product to any public waters. Permits may be required to treat such water.

Recreational Use of Water in Treated Areas: There are no restrictions on the use of water in treated areas for recreational purposes, including swimming and fishing.

Livestock Use of Water in/from Treated Areas:

There are no restrictions on livestock consumption of water from treated areas.

Precautions for Potable Water Intakes: DO NOT apply Alligare Ecomazapyr 2 SL directly to waters within one-half mile upstream of an active potable water intake in flowing water (i.e., river, stream, etc.) or within one-half mile of an active potable water intake in a standing body of water such as lake, pond or reservoir. To make aquatic applications around and within one-half mile of active potable water intakes, the water intake must be turned off during application and for at least 48 hours after the treatment. Such aquatic applications may be made only where there are alternative water sources or holding ponds that can be used while active potable water intakes are turned off during the minimum 48 hours post-treatment period. Note: Existing potable water intakes which are no longer in use, such as those replaced by connections to wells or municipal water systems, are not considered to be active potable water intakes. This restriction does not apply to intermittent, inadvertent overspray of water in terrestrial use sites.

APPLICATION TO WATERS USED FOR IRRIGATION

DO NOT use water treated with Alligare Ecomazapyr 2 SL for irrigation purposes for 120 days after treatment or until Alligare Ecomazapyr 2 SL residue levels are determined by laboratory analysis, or other appropriate means of analysis, to be 1.0 ppb or less.

Seasonal Irrigation Waters: Alligare Ecomazapyr 2 SL may be applied during the off-season to surface waters that are used for irrigation on a seasonable basis, provided at least 120 days elapses between Alligare Ecomazapyr 2 SL treatment and the first use of treated water for irrigation or until Alligare Ecomazapyr 2 SL residue levels are determined by laboratory analysis, or other appropriate means of analysis, to be 1.0 ppb or less.

Irrigation Canals/Ditches: DO NOT apply Alligare Ecomazapyr 2 SL to irrigation canals/ditches unless the 120-day restriction on irrigation water usage can be observed or Alligare Ecomazapyr 2 SL residue levels are determined by laboratory analysis, or other appropriate means of analysis, to be 1.0 ppb or less. DO NOT apply Alligare Ecomazapyr 2 SL to dry irrigation canals/ditches.

Quiescent or Slow Moving Waters: In lakes and reservoirs DO NOT apply Alligare Ecomazapyr 2 SL within one (1) mile of an active irrigation water intake during the irrigation season. Applications less than one (1) mile from an inactive irrigation water intake may be made during the off-season, provided the irrigation intake will remain inactive for at least 120 days after treatment or until Alligare Ecomazapyr 2 SL residue levels are determined by laboratory analysis, or other appropriate means of analysis, to be 1.0 ppb or less.

Moving Waters: DO NOT apply Alligare Ecomazapyr 2 SL within one-half mile downstream of an active irrigation water intake. When making applications upstream from an active irrigation water intake, the intake must be turned off for a period of time sufficient to allow the upstream treated water to completely flow past the irrigation intake. Shut off time will be determined by the speed of water flow and the distance and length of water treated upstream from the intake. Consult local, state and/or federal authorities before making any applications upstream from an active irrigation water intake.

TANK MIXES

Alligare Ecomazapyr 2 SL may be tank-mixed with other aquatic-use herbicides to broaden the spectrum of control of emergent and floating aquatic vegetation provided that the tank mix product label does not prohibit such mixing. Consult manufacturer's labels for specific rates and weeds controlled. Always follow the more restrictive label when making an application involving tank-mixes.

AERIAL APPLICATIONS

Exercise all precautions to minimize or eliminate spray drift. Aerial applications may only be made by helicopter; applications may not be made by airplane. Helicopters can be used to apply **Alligare Ecomazapyr 2 SL**; however, **DO NOT** apply by helicopter unless appropriate buffer zones can be maintained to prevent spray drift out of the target area or where damage to desirable vegetation can be tolerated. Helicopters equipped with a Microfoil™ boom, Thru-Valve™ boom or raindrop nozzles must be used and calibrated. Unless applying with a Microfoil boom, use a drift control agent at the recommended label rate. To avoid drift, **DO NOT** make applications during inversion conditions, when winds are gusty, or during any other conditions that promote spray drift. Side trimming is not recommended with **Alligare Ecomazapyr 2 SL** unless death of treated vegetation is acceptable.

Uniformly apply **Alligare Ecomazapyr 2 SL** in 5 to 30 gallons of water per acre. Use a nonionic surfactant, methylated seed oil or silicone-based surfactant (see the **ADJUVANTS** section of this label for specific recommendations). An anti-foam agent may be added, if needed.

Thoroughly clean application equipment, including landing gear, by thoroughly flushing with water immediately after using this product. Prolonged exposure of uncoated/unpainted steel (except stainless steel) surfaces to this product may result in corrosion and failure of the exposed part. Maintaining painted surfaces may prevent corrosion.

GROUND APPLICATIONS

FOLIAR APPLICATIONS:

For Low Volume Foliar Applications: Use equipment calibrated to deliver 5 to 20 gallons of spray solution per acre. Thoroughly mix 0.5 to 5% (v/v) **Alligare Ecomazapyr 2 SL** in water plus surfactant (see the **ADJUVANTS** section of this label for recommendations). Use an anti-foam agent at the recommended rate, if needed. For difficult to control species (see **AQUATIC WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL** section and the **ADDITIONAL WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL** section for relative susceptibility of weed species), apply the higher concentrations of herbicide and/or spray volumes but **DO NOT** apply more than 6 pints of **Alligare Ecomazapyr 2 SL** per acre. Excessive wetting of foliage is not recommended. See the **SPRAY SOLUTION MIXING GUIDE** below for some suggested volumes of **Alligare Ecomazapyr 2 SL** and water.

Select appropriate nozzles to avoid over-application. Proper application is critical to ensure desirable results. Optimum results are achieved when the spray covers the crown and approximately 70 percent of the plant. The use of a flat fan nozzle tip with a spray angle of 40 degrees or less will aid in proper deposition.

Recommended nozzle tip sizes include 4004E or 1504E. For a straight stream and cone pattern, use adjustable cone nozzles such as 5500 X3 or 5500 X4. Attaching a rollover valve onto a Spraying Systems Model 30 gunjet or other similar spray guns allows for the use of both a flat fan and cone tips on the same gun.

Moisten, but do not drench target vegetation. **DO NOT** spray to run off.

For Low Volume Foliar Applications by Backpacks: For low-growing species, spray downward to cover approximately 70% of the plant and the crown.

For target species 4 to 8 feet tall, apply a directed spray in a smooth vertical motion from the crown upward on at least two sides of the target vegetation, making sure to cover the crown whenever possible.

For target species over 8 feet tall, apply a directed spray in a smooth zig-zag motion from the crown upward on at least two sides of the target brush.

For Low Volume Foliar Applications with Hydraulic Handgun Equipment: Use same technique as described above for **Low Volume with Backpacks**.

For broadcast applications, simulate a gentle rain near the top of target vegetation, allowing spray to penetrate the target foliage and contact the crown without run-off onto understory vegetation. **DO NOT** spray to run-off. Herbicide spray that contacts understory vegetation may result in severe injury or death of understory plants.

SPRAY SOLUTION MIXING GUIDE FOR LOW-VOLUME FOLIAR APPLICATIONS

Amount of spray solution	Desired Concentration (fluid volume)				
	0.5%	0.75%	1%	1.5%	5%
(amount of Alligare Ecomazapyr 2 SL to use)					
1 gallon	0.6 oz	0.9 oz	1.3 oz	1.9 oz	6.5 oz
3 gallons	1.9 oz	2.8 oz	3.8 oz	5.8 oz	1.2 pint
4 gallons	2.5 oz	3.8 oz	5.1 oz	7.7 oz	1.6 pint
5 gallons	3.2 oz	4.8 oz	6.5 oz	9.6 oz	2 pints
50 gallons	2 pints	3 pints	4 pints	6 pints	10 pints
100 gallons	4 pints	6 pints	8 pints	6 quarts	5 gallons

2 tablespoons = 1 fluid ounce

For High Volume Foliar Applications: For optimum performance when spraying medium to high-density vegetation, use equipment calibrated to deliver up to 100 gallons of finished spray per acre (GPA). Application volumes exceeding 100 gallons GPA may result in excessive spray run-off, causing injury to desirable ground cover species. Thoroughly mix **Alligare Ecomazapyr 2 SL** in water and include a surfactant (see **ADJUVANTS** section for surfactant recommendations). Use an anti-foam agent according to its label, if needed. For hard-to-control species (see **AQUATIC WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL** section and the **ADDITIONAL WEEDS CONTROLLED BY ALLIGARE ECOMAZAPYR 2 SL** section for relative susceptibility of weeds), use the higher concentrations of the herbicide and/or spray volumes, but **DO NOT** apply more than 6 pints of **Alligare Ecomazapyr 2 SL** per acre. Uniformly cover the foliage of the target vegetation but **DO NOT** apply to run-off.

Side Trimming: **DO NOT** side trim with **Alligare Ecomazapyr 2 SL** unless severe injury or death of the treated vegetation is acceptable. **Alligare Ecomazapyr 2 SL** is readily translocated and can result in the death of the vegetation.

CUT SURFACE TREATMENTS

Alligare Ecomazapyr 2 SL will control undesirable woody vegetation when applied as a water solution to the cambium area of freshly-cut stump surfaces or to cuts on the stem of the target woody vegetation. Applications can be made at any time of the year except during periods of heavy sap flow in the spring. Tree injection and cut stem treatments are most effective in late summer and early fall. **DO NOT** over apply solution causing run-off from the cut surface.

NOTE: Injury may occur to desirable woody plants if the shoots extend from the same root system or their root systems are grafted to those of the treated tree.

CUT SURFACE APPLICATIONS WITH DILUTE AND CONCENTRATE SOLUTIONS:

Mix **Alligare Ecomazapyr 2 SL** as either a concentrated or dilute solution for cut surface treatments. Apply dilute solutions to the cut surface of the stump or to cuts on the stem of the target woody vegetation. Apply concentrate solutions to cuts on the stem. Use of concentrated solutions permits application to fewer cuts on the stem, especially for large diameter trees. Follow the application directions below to determine proper application techniques for each type of solution.

To prepare a dilute solution, mix 8 to 12 fluid ounces of **Alligare Ecomazapyr 2 SL** with one gallon of water. If temperatures are such that freezing of the spray mixture may occur, antifreeze (ethylene glycol) may be added according to manufacturer's label to prevent freezing. The use of a surfactant or penetrating agent may improve herbicide uptake through partially callused cambium tissues.

To prepare a concentrated solution, mix 2 quarts of **Alligare Ecomazapyr 2 SL** with no more than 1 quart water.

For cut stump treatments: Spray or brush the solution onto the cambium area of the freshly cut stump surface. Thoroughly wet the entire cambium area (the wood just inside the bark of the stump).

For cut stem (injection, hack and squirt) treatments:

Dilute Solutions: Using standard injection equipment, apply 1 milliliter of solution at each injection site around the tree with no more than one-inch intervals between cut edges. Insure that the injector completely penetrates the bark at each injection site.

Concentrate Solutions: Using standard injection equipment, apply 1 milliliter of solution at each injection site. Make at least one injection cut for every 3 inches of Diameter at Breast Height (DBH) on the target tree. For example, a 3-inch DBH tree will receive 1 injection cut while a 6-inch DBH tree will receive 2 injection cuts. On trees requiring more than one injection site, place the injection cuts at approximately equal intervals around the tree.

Frill or girdle treatments: Use a hatchet, machete or chainsaw to make cuts through the bark around the tree to expose the cambium. The cuts should angle downward into the cambium sufficiently deep to expose at least two growth rings. Use a spray applicator or brush to apply a 25% to 100% solution of **Alligare Ecomazapyr 2 SL** into each cut until thoroughly wet. Avoid applying so much herbicide that runoff to the ground or water occurs.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: **DO NOT** store below 10°F.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Nonrefillable container (2.5, 30, 265 gallons): Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying.

(Nonrefillable \leq 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

(Nonrefillable $>$ 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Refillable container (15 gallon): Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

IMPORTANT: Read the entire **DIRECTIONS FOR USE** and the **CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY** before using this product. If terms are not acceptable, return the unopened product container at once.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

To the extent consistent with applicable law, upon purchase or use of this product, purchaser and user agree to the following terms:

Warranty: Alligare, LLC (the Company) warrants that this product conforms to the chemical description on the label in all material respects and is reasonably fit for the purpose referred to in the directions for use, subject to the exceptions noted below, which are beyond the Company's control. To the extent consistent with applicable law, the Company makes no other representation or warranty, express or implied, concerning the product, including no implied warranty of merchantability or fitness for a particular purpose. No such warranty shall be implied by law, and no agent or representative is authorized to make any such warranty on the Company's behalf.

Terms of Sale: The Company's directions for use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, and the manner of use or application (including failure to adhere to label directions), all of which are beyond the Company's control. To the extent consistent with applicable law, all such risks are assumed by the user.

Limitation of Liability: To the extent consistent with applicable law, the exclusive remedy against the Company for any cause of action relating to the handling or use of this product is a claim for damages, and in no event shall damages or any other recovery of any kind exceed the price of the product which caused the alleged loss, damage, injury or other claim. To the extent consistent with applicable law, under no circumstances shall the Company be liable for any special, indirect, incidental or consequential damages of any kind, including loss of profits or income, and any such claims are hereby waived. Some states do not allow the exclusion or limitation of incidental or consequential damages.

The Company and the seller offer this product, and the purchaser and user accept this product, subject to the foregoing warranty, terms of sale and limitation of liability, which may be varied or modified only by an agreement in writing signed on behalf of the Company by an authorized representative.

Microfoil is a trademark of Rhone Poulenc Ag. Company.
Thru-Valve is a trademark of Waldrum Specialties.

EPA 20080825

PYXIS REGULATORY CONSULTING, INC.

4110 136th St. NW
Gig Harbor, WA 98332

Phone: 253-853-7369
Fax: 253-853-5516
www.PyxisRC.com

August 25, 2008

COURIER DELIVERY

Jim Tompkins (PM 25)
Document Processing Desk (**NOTIF**)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 S. Crystal Drive
Arlington, VA 22202

RE: Alligare, LLC – Alligare Ecomazapyr 2 SL (EPA Reg. No. 81927-22)
Revision to Container Disposal Instructions per PRN 2007-4

Dear Mr. Tompkins,


On behalf of Alligare, LLC please find the enclosed label notification revising the container disposal instructions per PRN 2007-4 for Alligare's Ecomazapyr 2 SL (EPA Reg. No. 81927-22).

In support of this notification submission, we submit the following documents:

1. Completed Application for Registration (EPA Form 8570-1)
2. One (1) copy of the Alligare Ecomazapyr 2 SL labeling with changes tracked
3. One (1) copy of the Alligare Ecomazapyr 2 SL labeling with changes incorporated
4. Letter of Authorization

Please feel free to contact me by phone (253) 853-7369 or by email at Mike@PyxisRC.com if you have any questions or need any additional information.


Sincerely,



Michael Kellogg

Enclosures

cc: Jay Golz; Alligare, LLC





PROTECTING:
RIGHT OF WAY
UTILITIES
RAILROAD
RANGE & PASTURELAND
FORESTRY
AQUATIC
INDUSTRIAL

June 3, 2008

To Whom It May Concern:

RE: Letter of Authorization

Dear Sir or Madam:

Please let this letter serve to confirm that Pyxis Regulatory Consulting, Inc. is authorized to act as agent for Alligare, LLC (EPA Company Number 81927), before the U.S. Environmental Protection Agency and state governmental agencies in all matters regarding our pesticide registrations pursuant to the Federal Insecticide, Fungicide and Rodenticide Act ("FIFRA"), 7 U.S.C. § 136 et seq. and state law.

If you have any questions, please do not hesitate to contact me.

Sincerely,

Jay Golz
Vice-President & COO

cc: Pyxis Regulatory Consulting, Inc.



United States
Environmental Protection Agency
Washington, DC 20460

☐ Registration
☐ Amendment
☒ Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 81927-22	2. EPA Product Manager J. Tompkins	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Alligare, LLC / Alligare Ecomazapyr 2 SL	PM# 25	
5. Name and Address of Applicant (Include ZIP Code) Alligare, LLC c/o Pyxis Regulatory Consulting, Inc. 4110 136th St. NW Gig Harbor, WA 98332 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. <u>SEP 11 2008</u> Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of label change per PR Notice 2007-4. This notification is consistent with the guidance in PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input checked="" type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt.	No. per container
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 2.5, 15, 30, 265 gallon		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____			

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			
Name Michael Kellogg	Title Agent	Telephone No. (Include Area Code) (253) 853-7369	
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.			6. Date Application Received (Stamped)
2. Signature 	3. Title Agent		
4. Typed Name Michael Kellogg	5. Date 8/25/08		